



Sub-riemannian geometry from intrinsic viewpoint

Marius Buliga

► **To cite this version:**

Marius Buliga. Sub-riemannian geometry from intrinsic viewpoint. École de recherche CIMPA : Géométrie sous-riemannienne, Jan 2012, BEYROUTH, Lebanon. <hal-00700925v2>

HAL Id: hal-00700925

<https://hal-confremo.archives-ouvertes.fr/hal-00700925v2>

Submitted on 19 Jun 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Sub-riemannian geometry from intrinsic viewpoint

Marius Buliga

Institute of Mathematics, Romanian Academy
P.O. BOX 1-764, RO 014700

București, Romania

Marius.Buliga@imar.ro

This version: 14.06.2012

Abstract

Gromov proposed to extract the (differential) geometric content of a sub-riemannian space exclusively from its Carnot-Carathéodory distance. One of the most striking features of a regular sub-riemannian space is that it has at any point a metric tangent space with the algebraic structure of a Carnot group, hence a homogeneous Lie group. Siebert characterizes homogeneous Lie groups as locally compact groups admitting a contracting and continuous one-parameter group of automorphisms. Siebert result has not a metric character.

In these notes I show that sub-riemannian geometry may be described by about 12 axioms, without using any a priori given differential structure, but using dilation structures instead. Dilation structures bring forth the other intrinsic ingredient, namely the dilations, thus blending Gromov metric point of view with Siebert algebraic one.

MSC2000: 51K10, 53C17, 53C23

1	Introduction	2
2	Metric spaces, groupoids, norms	4
2.1	Normed groups and normed groupoids	5
2.2	Gromov-Hausdorff distance	7
2.3	Length in metric spaces	8
2.4	Metric profiles. Metric tangent space	10
2.5	Curvdimension and curvature	12
3	Groups with dilations	13
3.1	Conical groups	14
3.2	Carnot groups	14
3.3	Contractible groups	15
4	Dilation structures	16
4.1	Normed groupoids with dilations	16
4.2	Dilation structures, definition	18

5	Examples of dilation structures	20
5.1	Snowflakes, nonstandard dilations in the plane	20
5.2	Normed groups with dilations	21
5.3	Riemannian manifolds	22
6	Length dilation structures	22
7	Properties of dilation structures	24
7.1	Metric profiles associated with dilation structures	24
7.2	The tangent bundle of a dilation structure	26
7.3	Differentiability with respect to a pair of dilation structures	29
7.4	Equivalent dilation structures	30
7.5	Distribution of a dilation structure	31
8	Supplementary properties of dilation structures	32
8.1	The Radon-Nikodym property	32
8.2	Radon-Nikodym property, representation of length, distributions	33
8.3	Tempered dilation structures	34
9	Dilation structures on sub-riemannian manifolds	37
9.1	Sub-riemannian manifolds	37
9.2	Sub-riemannian dilation structures associated to normal frames	38
10	Coherent projections: a dilation structure looks down on another	41
10.1	Coherent projections	42
10.2	Length functionals associated to coherent projections	44
10.3	Conditions (A) and (B)	45
11	Distributions in sub-riemannian spaces as coherent projections	45
12	An intrinsic description of sub-riemannian geometry	47
12.1	The generalized Chow condition	47
12.2	The candidate tangent space	50
12.3	Coherent projections induce length dilation structures	53

1 Introduction

In these notes I show that sub-riemannian geometry may be described intrinsically, in terms of dilation structures, **without using any a priori given differential structure**.

A complete riemannian manifold is a length metric space by the Hopf-Rinow theorem. The problem of intrinsic characterization of riemannian spaces asks for the recovery of the manifold structure and of the riemannian metric from the distance function (associated to the length functional).

For 2-dim riemannian manifolds the problem has been solved by A. Wald in 1935 [34]. In 1948 A.D. Alexandrov [1] introduces its famous curvature (which uses comparison triangles) and proves that, under mild smoothness conditions on this curvature, one is capable to recover the differential structure and the metric of the 2-dim riemannian manifold. In 1982 Alexandrov proposes as a conjecture that a characterization of a riemannian manifold (of any dimension) is possible in terms of metric (sectional) curvatures (of the type introduced by Alexandrov) and weak smoothness assumptions formulated in metric way (as for example Hölder smoothness).

The problem has been solved by Nikolaev [28] in 1998. He proves that every locally compact length metric space M , not linear at one of its points, with α -Hölder continuous metric sectional curvature of the generalized tangent bundle $T^m(M)$ (for some $m = 1, 2, \dots$, which admits local geodesic extendability, is isometric to a C^{m+2} smooth riemannian manifold. We shall remain vague about what is the meaning of: not linear, metric sectional curvature, generalized tangent bundle. Please read the excellent paper by Nikolaev for grasping the precise meaning of the result. Nevertheless, we may summarize the solution of Nikolaev like this:

- He constructs a (family of) intrinsically defined tangent bundle(s) of the metric space, by using a generalization of the cosine formula for estimating a kind of a distance between two curves emanating from different points. This will lead him to a generalization of the tangent bundle of a riemannian manifold endowed with the canonical Sasaki metric.
- He defines a notion of sectional curvature at a point of the metric space, as a limit of a function of nondegenerated geodesic triangles, limit taken as these triangles converge (in a precised sense) to the point.
- The sectional curvature function thus constructed is supposed to satisfy a smoothness condition formulated in metric terms.

He proves that under the smoothness hypothesis he gets the conclusion.

In this paper we prove (see theorem 8.10):

Theorem 1.1 *The dilation structure associated to a riemannian manifold, as in proposition 5.2, is tempered (definition 8.6), has the Radon-Nikodym property (definition 8.1) and is a length dilation structure (definition 6.3).*

If (X, d, δ) is a strong dilation structure which is tempered, it has the Radon-Nikodym property and moreover for any $x \in X$ the tangent space in the sense of dilation structures (definition 7.4) is a commutative local group, then any open, with compact closure subset of X can be endowed with a C^1 riemannian structure which gives a distance d' which is bilipschitz equivalent with d .

Sub-riemannian spaces are length metric spaces as well, why are them different? First of all, any riemannian space is a sub-riemannian one, therefore sub-riemannian spaces are more general than riemannian ones. It is not clear at first sight why the characterization of riemannian spaces does not extend to sub-riemannian ones. In fact, there are two problematic steps for such a program for extending Nikolaev result to sub-riemannian spaces: the cosine formula, as well as the Sasaki metric on the tangent bundle don't have a correspondent in sub-riemannian geometry (there is, basically, no statement canonically corresponding to Pythagoras theorem); the sectional curvature at a point cannot be introduced by means of comparison triangles, because sub-riemannian spaces do not behave well with respect to this comparison of triangle idea.

The problem of intrinsic characterization of sub-riemannian spaces has been formulated by Gromov in [25]. Gromov takes the Carnot-Carathéodory distance as the only intrinsic object of a sub-riemannian space. Indeed, in [25]. section 0.2.B. he writes:

"If we live inside a Carnot-Carathéodory metric space V we may know nothing whatsoever about the (external) infinitesimal structures (i.e. the smooth structure on V , the subbundle $H \subset T(V)$ and the metric g on H) which were involved in the construction of the CC metric."

He then formulates two main problems:

- (1) *"Develop a sufficiently rich and robust internal C-C language which would enable us to capture the essential external characteristics of our C-C spaces"*. (he proposes as an example to recognize the rank of the horizontal distribution, but in my opinion this is, say, something much less essential than to "recognize" the "differential structure", in the sense proposed here as the equivalence class under local equivalence of dilation structures, see definition 7.9)

(2) "Develop external (analytic) techniques for evaluation of internal invariants of V ."

Especially the problem (2) raises a big question: what is, in fact, a sub-riemannian space? Should it be defined only in relation with the differential geometric construction using horizontal distributions and CC distance? In this paper we propose that a sub-riemannian space is a particular case of a pair of spaces, one looking down on another. The "sub-riemannian"-ness is a relative notion.

As for the problem (1), a solution is proposed here, by using dilation structure. The starting point is to remark that a regular sub-riemannian space has metric tangent spaces with the structure of a Carnot group. All known proofs are using indeed the intrinsic CC distance, but also the differential structure of the manifold and the differential geometric definition of the CC distance. The latter are not intrinsic, according to Gromov criterion, even if the conclusion (the tangent space has an algebraic Carnot group structure) looks intrinsic. But is this Carnot group structure intrinsic or is an artifact of the method of proof which was used?

Independently, there is an interest into the characterization of contractible topological groups. A result of Siebert [30] characterizes homogeneous Lie groups as locally compact groups admitting a contracting and continuous one-parameter group of automorphisms. This result is relevant because, we argued before, the Carnot group structure comes from the self-similar metric structure of a tangent space, via the result of Siebert.

If we enlarge the meaning of "intrinsic", such as to contain the CC distance, but also the approximate self-similar structure of the sub-riemannian space, then we are able to give a characterization of these spaces. This approximate self-similarity is modeled by dilation structures (initially called "dilatation structures" [4], a nod to the latin origin of the word). Dilation structures bring forth the other intrinsic ingredient, namely the dilations, which are generalizations of Siebert' contracting group of automorphisms.

According to the characterization given in this paper, a regular sub-riemannian space is one which can be constructed from: a tempered dilation structure with the Radon-Nikodym property and commutative tangent spaces, and from a coherent projection which satisfies (Cgen), (A) and (B) properties. As a corollary, we recover the known fact that sub-riemannian spaces have the Radon-Nikodym property and we learn the new fact that they are length dilation structures, which provides a characterization of the behaviour of the rescaled length functionals induced by the Carnot-Carathéodory distance.

As it is maybe to be expected from a course notes paper, these notes are based on previous papers of mine, mainly [7] (section 12 follows almost verbatim the section 10 of [7]), [4], [5], [6], [9] and also from a number of arxiv papers of mine, mentioned in the bibliography. Many clarifications and theorems are added, in order to construct over the foundations laid elsewhere. I hope that the unitary presentation will help the understanding of the subject.

Acknowledgements. These are the notes prepared for the course "Metric spaces with dilations and sub-riemannian geometry from intrinsic point of view", CIMPA research school on sub-riemannian geometry (2012). Unfortunately I have not been able to attend the school. I want to express my thanks to the organizers for inviting me and also my excuses for not being there. This work was partially supported by a grant of the Romanian National Authority for Scientific Research, CNCS UEFISCDI, project number PN-II-ID-PCE-2011-3-0383.

2 Metric spaces, groupoids, norms

Metric spaces have been introduced by Fréchet (1906) in the paper [22].

Definition 2.1 A metric space (X, d) is a pair formed by a set X and a function called distance, $d : X \times X \rightarrow [0, +\infty)$, which satisfies the following: (i) $d(x, y) = 0$ if and only if $x = y$; (ii)

(symmetry) for any $x, y \in X$ we have $d(x, y) = d(y, x)$; (iii) (triangle inequality) for any $x, y, z \in X$ we have $d(x, z) \leq d(x, y) + d(y, z)$. The ball of radius $r > 0$ and center $x \in X$ is the set $B(x, r) = \{y \in X : d(x, y) < r\}$. Sometimes we may use the notation $B_d(x, r)$, which indicates the dependence on the distance d .

The topology and uniformity on the metric space (X, d) is the one generated by balls, respectively by preimages of the distance fonction.

2.1 Normed groups and normed groupoids

Starting from the observation that normed trivial groupoids are in bijective correspondence with metric spaces, proposition 2.5, we think it is interesting to extend the theory of metric spaces to normed groupoids. This is explained in detail in [14].

Groups are groupoids with one object, so let us start with normed groups, then pass to normed groupoids.

Definition 2.2 A normed group is a pair (G, ρ) , formed by a group G , with operation $(x, y) \in G \times G \mapsto xy$, the inverse $x \in G \mapsto x^{-1}$ and neutral element denoted by e , and a norm $\rho : G \rightarrow [0, +\infty)$ which satisfies the following: (i) $\rho(x) = 0$ if and only if $x = e$; (ii) (symmetry) for any $x \in G$ we have $\rho(x^{-1}) = \rho(x)$; (iii) (sub-additivity) for any $x, y \in G$ we have $\rho(xy) \leq \rho(x) + \rho(y)$.

A normed group (G, ρ) can be seen as a metric space. Indeed, as expected, the norm ρ induce distances, left or right invariant:

$$d_L(x, y) = \rho(x^{-1}y) \quad , \quad d_R(x, y) = \rho(xy^{-1}) \quad .$$

Groupoids are generalization of groups. We shall model them by looking to the set of arrows, which is a set with a partially defined binary operation and a unary operation (the inverse function). A groupoid norm will be a function defined on the set of arrows, with properties similar with the ones of a norm over a group.

Definition 2.3 A normed groupoid (G, ρ) is a pair formed by:

- a groupoid G , i.e. a set with two partially defined operations: the composition $m : G^{(2)} \subset G \times G \rightarrow G$, denoted multiplicatively $m(a, b) = ab$, and the inverse $inv : G \rightarrow G$, denoted $inv(a) = a^{-1}$. The operations satisfy: for any $a, b, c \in G$

- (i) if $(a, b) \in G^{(2)}$ and $(b, c) \in G^{(2)}$ then $(a, bc) \in G^{(2)}$ and $(ab, c) \in G^{(2)}$ and we have $a(bc) = (ab)c$,
- (ii) $(a, a^{-1}) \in G^{(2)}$ and $(a^{-1}, a) \in G^{(2)}$,
- (iii) if $(a, b) \in G^{(2)}$ then $abb^{-1} = a$ and $a^{-1}ab = b$.

The set of objects of the groupoid $X = Ob(G)$ is formed by all products $a^{-1}a$, $a \in G$. For any $a \in G$ we let $\alpha(a) = a^{-1}a \in X$ to be the source (object) of a and $\omega(a) = aa^{-1} \in X$ to be the target of a .

- a (groupoid) norm $d : G \rightarrow [0, +\infty)$ which satisfies:

- (i) $d(g) = 0$ if and only if $g \in Ob(G)$,
- (ii) (symmetry) for any $g \in G$, $d(g^{-1}) = d(g)$,
- (iii) (sub-additivity) for any $(g, h) \in G^{(2)}$, $d(gh) \leq d(g) + d(h)$,

If $Ob(G)$ is a singleton then G is just a group and the previous definition corresponds exactly to the definition 2.2 of a normed group. As in the case of normed groups, normed groupoids induce metric spaces too.

Proposition 2.4 *Let (G, d) be a normed groupoid. For any $x \in Ob(G)$ the pair $(\alpha^{-1}(x), d_x)$ is a metric space, where for any $g, h \in G$ with $\alpha(g) = \alpha(h) = x$ we define $d_x(g, h) = d(gh^{-1})$.*

Moreover, any normed groupoid is a disjoint union of metric spaces

$$G = \bigcup_{x \in Ob(G)} \alpha^{-1}(x) \quad , \quad (1)$$

such that for any $u \in G$ the "right translation"

$$R_u : \alpha^{-1}(\omega(u)) \rightarrow \alpha^{-1}(\alpha(u)) \quad , \quad R_u(g) = gu$$

is an isometry, that is for any $g, h \in \alpha^{-1}(\omega(u))$

$$d_{\omega(u)}(g, h) = d_{\alpha(u)}(R_u(g), R_u(h)) \quad .$$

Proof. If $\alpha(g) = \alpha(h) = x$ then $(g, h^{-1}) \in G^{(2)}$, therefore $d_x(g, h)$ is well defined. The proof of the first part is straightforward, the properties (i), (ii), (iii) of the groupoid norm ρ transform respectively into (i), (ii), (iii) properties of the distance d_x .

For the second part of the proposition remark that R_u is well defined and moreover $R_u(g)(R_u(h))^{-1} = gh^{-1}$. Then

$$\begin{aligned} d_{\alpha(u)}(R_u(g), R_u(h)) &= d\left(R_u(g)(R_u(h))^{-1}\right) = \\ &= d(gh^{-1}) = d_{\omega(u)}(g, h) \end{aligned}$$

and the proof is done. \square

Conversely, any metric space is identified with a normed groupoid.

Proposition 2.5 *Let (X, d) be a metric space and consider the "trivial groupoid" $G = X \times X$, with multiplication $(x, y)(y, z) = (x, z)$ and inverse $(x, y)^{-1} = (y, x)$. Then (G, d) is a normed groupoid and moreover any component of the decomposition (1) of G is isometric with (X, d) .*

Moreover if $G = X \times X$ is the trivial groupoid associated to the set X and d is a norm on G then (X, d) is a metric space.

Proof. We have $\alpha(x, y) = (y, y)$ and $\omega(x, y) = (x, x)$, therefore the set of objects of the trivial groupoid is $Ob(G) = \{(x, x) : x \in X\}$. This set can be identified with X by the bijection $(x, x) \mapsto x$. Moreover, for any $x \in X$ we have $\alpha^{-1}((x, x)) = X \times \{x\}$.

The distance $d : X \times X \rightarrow [0, +\infty)$ is a groupoid norm, seen as $d : G \rightarrow [0, +\infty)$. Indeed (i) $(d(x, y) = 0$ if and only if $(x, y) \in Ob(G)$) is equivalent with $d(x, y) = 0$ if and only if $x = y$. The symmetry condition (ii) is just the symmetry of the distance $d(x, y) = d(y, x)$. Finally the sub-additivity of d as a groupoid norm is equivalent with the triangle inequality. In conclusion $(X \times X, d)$ is a normed groupoid if and only if (X, d) is a metric space.

For any $x \in X$, let us compute the distance $d_{(x, x)}$, which is the distance on the space $\alpha^{-1}((x, x))$. We have

$$d_{(x, x)}((u, x), (v, x)) = d((u, x)(v, x)^{-1}) = d((u, x)(x, v)) = d(u, v)$$

therefore the metric space $(\alpha^{-1}((x, x)), d_{(x, x)})$ is isometric with (X, d) by the isometry $(u, x) \mapsto u$, for any $u \in X$. \square

In conclusion normed groups make good examples of metric spaces.

2.2 Gromov-Hausdorff distance

We shall denote by CMS the set of isometry classes of compact metric spaces. This set is endowed with the Gromov distance and with the topology is induced by this distance.

The Gromov-Hausdorff distance shall be introduced by way of (cartographic like) maps. Although this definition is well known, the cartographic analogy was explained first time in detail in [15], which we follow here.

Definition 2.6 *Let $\rho \subset X \times Y$ be a relation which represents a map of $dom \rho \subset (X, d)$ into $im \rho \subset (Y, D)$. To this map are associated three quantities: accuracy, precision and resolution.*

The accuracy of the map ρ is defined by:

$$acc(\rho) = \sup \{ | D(y_1, y_2) - d(x_1, x_2) | : (x_1, y_1) \in \rho, (x_2, y_2) \in \rho \} \quad (2)$$

The resolution of ρ at $y \in im \rho$ is

$$res(\rho)(y) = \sup \{ d(x_1, x_2) : (x_1, y) \in \rho, (x_2, y) \in \rho \} \quad (3)$$

and the resolution of ρ is given by:

$$res(\rho) = \sup \{ res(\rho)(y) : y \in im \rho \} \quad (4)$$

The precision of ρ at $x \in dom \rho$ is

$$prec(\rho)(x) = \sup \{ D(y_1, y_2) : (x, y_1) \in \rho, (x, y_2) \in \rho \} \quad (5)$$

and the precision of ρ is given by:

$$prec(\rho) = \sup \{ prec(\rho)(x) : x \in dom \rho \} \quad (6)$$

We may need to perform also a "cartographic generalization", starting from a relation ρ , with domain $M = dom(\rho)$ which is ε -dense in (X, d) .

Definition 2.7 *A subset $M \subset X$ of a metric space (X, d) is ε -dense in X if for any $u \in X$ there is $x \in M$ such that $d(x, u) \leq \varepsilon$.*

Let $\rho \subset X \times Y$ be a relation such that $dom \rho$ is ε -dense in (X, d) and $im \rho$ is μ -dense in (Y, D) . We define then $\bar{\rho} \subset X \times Y$ by: $(x, y) \in \bar{\rho}$ if there is $(x', y') \in \rho$ such that $d(x, x') \leq \varepsilon$ and $D(y, y') \leq \mu$.

If ρ is a relation as described in definition 2.7 then accuracy $acc(\rho)$, ε and μ control the precision $prec(\rho)$ and resolution $res(\rho)$. Moreover, the accuracy, precision and resolution of $\bar{\rho}$ are controlled by those of ρ and ε, μ , as well.

Proposition 2.8 *Let ρ and $\bar{\rho}$ be as described in definition 2.7. Then:*

- (a) $res(\rho) \leq acc(\rho)$,
- (b) $prec(\rho) \leq acc(\rho)$,
- (c) $res(\rho) + 2\varepsilon \leq res(\bar{\rho}) \leq acc(\rho) + 2(\varepsilon + \mu)$,
- (d) $prec(\rho) + 2\mu \leq prec(\bar{\rho}) \leq acc(\rho) + 2(\varepsilon + \mu)$,
- (e) $| acc(\bar{\rho}) - acc(\rho) | \leq 2(\varepsilon + \mu)$.

Proof. Remark that (a), (b) are immediate consequences of definition 2.6 and that (c) and (d) must have identical proofs, just by switching ε with μ and X with Y respectively. I shall prove therefore (c) and (e).

For proving (c), consider $y \in Y$. By definition of $\bar{\rho}$ we write

$$\{x \in X : (x, y) \in \bar{\rho}\} = \bigcup_{(x', y') \in \rho, y' \in \bar{B}(y, \mu)} \bar{B}(x', \varepsilon)$$

Therefore we get

$$res(\bar{\rho})(y) \geq 2\varepsilon + \sup \{res(\rho)(y') : y' \in im(\rho) \cap \bar{B}(y, \mu)\}$$

By taking the supremum over all $y \in Y$ we obtain the inequality

$$res(\rho) + 2\varepsilon \leq res(\bar{\rho})$$

For the other inequality, let us consider $(x_1, y), (x_2, y) \in \bar{\rho}$ and $(x'_1, y'_1), (x'_2, y'_2) \in \rho$ such that $d(x_1, x'_1) \leq \varepsilon, d(x_2, x'_2) \leq \varepsilon, D(y'_1, y) \leq \mu, D(y'_2, y) \leq \mu$. Then:

$$d(x_1, x_2) \leq 2\varepsilon + d(x'_1, x'_2) \leq 2\varepsilon + acc(\rho) + d(y'_1, y'_2) \leq 2(\varepsilon + \mu) + acc(\rho)$$

Take now a supremum and arrive to the desired inequality.

For the proof of (e) let us consider for $i = 1, 2$ $(x_i, y_i) \in \bar{\rho}, (x'_i, y'_i) \in \rho$ such that $d(x_i, x'_i) \leq \varepsilon, D(y_i, y'_i) \leq \mu$. It is then enough to take absolute values and transform the following equality

$$\begin{aligned} d(x_1, x_2) - D(y_1, y_2) &= d(x_1, x_2) - d(x'_1, x'_2) + d(x'_1, x'_2) - D(y'_1, y'_2) + \\ &\quad + D(y'_1, y'_2) - D(y_1, y_2) \end{aligned}$$

into well chosen, but straightforward, inequalities. \square

The Gromov-Hausdorff distance is simply the optimal lower bound for the accuracy of maps of (X, d) into (Y, D) .

Definition 2.9 *Let $(X, d), (Y, D)$, be a pair of metric spaces and $\mu > 0$. We shall say that μ is admissible if there is a relation $\rho \subset X \times Y$ such that $dom \rho = X, im \rho = Y$, and $acc(\rho) \leq \mu$. The Gromov-Hausdorff distance between (X, d) and (Y, D) is the infimum of admissible numbers μ .*

This is a true distance on the set of isometry classes of pointed compact metric spaces.

2.3 Length in metric spaces

In a metric space, the distance function associates a number to a pair of points. In length metric spaces we have a length functional defined over Lipschitz curves. This functional is defined over a space of curves, therefore is a more sophisticated object. Length dilation structures are the correspondent of dilation structures for length metric spaces. It is surprising that, as far as I know, before [7] there were no previous efforts to describe the behaviour of the length functional restricted to smaller and smaller regions around a point in a length metric space.

The following definitions and results are standard, see for example the first chapter of [2].

Definition 2.10 *The (upper) dilation of a map $f : (X, d) \rightarrow (Y, D)$, in a point $u \in Y$ is*

$$Lip(f)(u) = \limsup_{\varepsilon \rightarrow 0} \sup \left\{ \frac{D(f(v), f(w))}{d(v, w)} : v \neq w, v, w \in B(u, \varepsilon) \right\}$$

Clearly, in the case of a derivable function $f : \mathbb{R} \rightarrow \mathbb{R}^n$ the upper dilation is $Lip(f)(t) = \|\dot{f}(t)\|$.

A function $f : (X, d) \rightarrow (Y, D)$ is Lipschitz if there is a positive constant C such that for any $x, y \in X$ we have $D(f(x), f(y)) \leq C d(x, y)$. The number $Lip(f)$ is the smallest such positive constant. Then for any $x \in X$ we have the obvious relation $Lip(f)(x) \leq Lip(f)$.

A curve is a continuous function $c : [a, b] \rightarrow X$. The image of a curve is called "path". Geometrically speaking, length measures paths, not curves, therefore the length functional, if defined over a class of curves, should be independent on the reparameterization of the path (image of the curve).

Definition 2.11 *Let (X, d) be a metric space. There are several ways to define a notion of length. The length of a curve with L^1 upper dilation $c : [a, b] \rightarrow X$ is $L(f) = \int_a^b Lip(c)(t) dt$. The variation of any curve $c : [a, b] \rightarrow X$ is*

$$Var(c) = \sup \left\{ \sum_{i=0}^n d(c(t_i), c(t_{i+1})) : a = t_0 < t_1 < \dots < t_n < t_{n+1} = b \right\}$$

The length of the path $A = c([a, b])$ is the one-dimensional Hausdorff measure of the path, i.e.

$$l(A) = \liminf_{\delta \rightarrow 0} \left\{ \sum_{i \in I} diam E_i : diam E_i < \delta, A \subset \bigcup_{i \in I} E_i \right\}$$

The definitions are not equivalent, but for any Lipschitz curve $c : [a, b] \rightarrow X$, we have

$$L(c) = Var(c) \geq \mathcal{H}^1(c([a, b]))$$

Moreover, if c is injective (i.e. a simple curve) then $\mathcal{H}^1(c([a, b])) = Var(f)$.

It is important to know the fact (which will be used repeatedly) that any Lipschitz curve c admits a reparametrisation c' such that $Lip(c')(t) = 1$ for almost any $t \in [a, b]$.

We associate a length functional to a metric space.

Definition 2.12 *We shall denote by l_d the length functional induced by the distance d , defined only on the family of Lipschitz curves. If the metric space (X, d) is connected by Lipschitz curves, then the length induces a new distance d_l , given by:*

$$d_l(x, y) = \inf \{ l_d(c([a, b])) : c : [a, b] \rightarrow X \text{ Lipschitz}, \\ c(a) = x, c(b) = y \}$$

A length metric space is a metric space (X, d) , connected by Lipschitz curves, such that $d = d_l$.

Lipschitz curves in complete length metric spaces are absolutely continuous. Indeed, here is the definition of an absolutely continuous curve (definition 1.1.1, chapter 1, [2]).

Definition 2.13 *Let (X, d) be a complete metric space. A curve $c : (a, b) \rightarrow X$ is absolutely continuous if there exists $m \in L^1((a, b))$ such that for any $a < s \leq t < b$ we have*

$$d(c(s), c(t)) \leq \int_s^t m(r) dr.$$

Such a function m is called a upper gradient of the curve c .

For a Lipschitz curve $c : [a, b] \rightarrow X$ in a complete length metric space such a function $m \in L^1((a, b))$ is the upper dilation $Lip(c)$. More can be said about the expression of the upper dilation. We need first to introduce the notion of metric derivative of a Lipschitz curve.

Definition 2.14 *A curve $c : (a, b) \rightarrow X$ is metrically derivable in $t \in (a, b)$ if the limit*

$$md(c)(t) = \lim_{s \rightarrow t} \frac{d(c(s), c(t))}{|s - t|}$$

exists and it is finite. In this case $md(c)(t)$ is called the metric derivative of c in t .

For the proof of the following theorem see [2], theorem 1.1.2, chapter 1.

Theorem 2.15 *Let (X, d) be a complete metric space and $c : (a, b) \rightarrow X$ be an absolutely continuous curve. Then c is metrically derivable for \mathcal{L}^1 -a.e. $t \in (a, b)$. Moreover the function $md(c)$ belongs to $L^1((a, b))$ and it is minimal in the following sense: $md(c)(t) \leq m(t)$ for \mathcal{L}^1 -a.e. $t \in (a, b)$, for each upper gradient m of the curve c .*

2.4 Metric profiles. Metric tangent space

To any locally compact metric space we associate a metric profile [11, 13]. This metric profile is a way of organizing the information given by the distance function in order to get an understanding of the local behaviour of the distance around a point of the space. We need to consider local compactness in order to have compact small balls in the next definition.

Let us denote by CMS' the set of isometry classes of pointed compact metric spaces. An element of CMS' is denoted like $[X, d, x]$ and is the equivalence class of a compact metric space (X, d) with a specified point $x \in X$, with respect to the following equivalence relation: two pointed compact metric spaces (X, d, x) and (Y, D, y) are equivalent if there is a surjective isometry $f : (X, d) \rightarrow (Y, D)$ such that $f(x) = y$.

The set CMS' is endowed with the GH distance for pointed metric spaces. In order to define this distance we have to slightly modify definition 2.9, by restricting the class of maps (relations) of (X, d) into $(Y, D) - \rho \subset X \times Y$ such that $dom \rho = X$, $im \rho = Y$ - to those which satisfy also $(x, y) \in \rho$.

We can define now metric profiles.

Definition 2.16 *The metric profile associated to the locally metric space (M, d) is the assignment (for small enough $\varepsilon > 0$)*

$$(\varepsilon > 0, x \in M) \mapsto \mathbb{P}^m(\varepsilon, x) = \left[\bar{B}(x, 1), \frac{1}{\varepsilon}d, x \right] \in CMS'$$

The metric profile of the space at a point is therefore a curve in another metric space, namely CMS' . with a Gromov-Hausdorff distance. It is not any curve, but one which has certain properties which can be expressed with the help of the GH distance. Indeed, for any $\varepsilon, b > 0$, sufficiently small, we have

$$\mathbb{P}^m(\varepsilon b, x) = \mathbb{P}_{d_b}^m(\varepsilon, x)$$

where $d_b = (1/b)d$ and $\mathbb{P}_{d_b}^m(\varepsilon, x) = [\bar{B}(x, 1), \frac{1}{\varepsilon}d_b, x]$.

These curves give interesting local and infinitesimal information about the metric space. For example, what kind of metric space has constant metric profile with respect to one of its points?

Definition 2.17 *A metric cone (X, d, x) is a locally compact metric space (X, d) , with a marked point $x \in X$ such that for any $a, b \in (0, 1]$ we have*

$$\mathbb{P}^m(a, x) = \mathbb{P}^m(b, x)$$

Metric cones are self-similar, in the sense that they have dilations.

Definition 2.18 Let (X, d, x) be a metric cone. For any $\varepsilon \in (0, 1]$ a dilation is a function $\delta_\varepsilon^x : \bar{B}(x, 1) \rightarrow \bar{B}(x, \varepsilon)$ such that

- $\delta_\varepsilon^x(x) = x$,
- for any $u, v \in X$ we have

$$d(\delta_\varepsilon^x(u), \delta_\varepsilon^x(v)) = \varepsilon d(u, v)$$

The existence of dilations for metric cones comes from the definition 2.17. Indeed, dilations are just isometries from $(\bar{B}(x, 1), d, x)$ to $(\bar{B}(x, \frac{1}{a}d, x)$.

Definition 2.19 A (locally compact) metric space (M, d) admits a (metric) tangent space in $x \in M$ if the associated metric profile $\varepsilon \mapsto \mathbb{P}^m(\varepsilon, x)$ admits a prolongation by continuity in $\varepsilon = 0$, i.e if the following limit exists:

$$[T_x M, d^x, x] = \lim_{\varepsilon \rightarrow 0} \mathbb{P}^m(\varepsilon, x) \quad (7)$$

The connection between metric cones, tangent spaces and metric profiles in the abstract sense is made by the following proposition.

Proposition 2.20 Metric tangent spaces are metric cones.

Proof. A tangent space $[V, d_v, v]$ exists if and only if we have the limit from the relation (7), that is of there exists a prolongation by continuity to $\varepsilon = 0$ of the metric profile $\mathbb{P}^m(\cdot, x)$. For any $a \in (0, 1]$ we have $\left[\bar{B}(x, 1), \frac{1}{a}d^x, x \right] = \lim_{\varepsilon \rightarrow 0} \mathbb{P}^m(a\varepsilon, x)$, therefore we have

$$\left[\bar{B}(x, 1), \frac{1}{a}d^x, x \right] = [T_x M, d^x, x]$$

which proves the thesis. \square

We may also define abstract metric profiles. The previously defined metric profiles are abstract metric profiles, but we shall see further (related to the Mitchell theorem in sub-riemannian geometry, for example) that abstract metric profiles are useful too.

Definition 2.21 An abstract metric profile is a curve $\mathbb{P} : [0, a] \rightarrow CMS$ such that

- (a) it is continuous at 0,
- (b) for any $b \in [0, a]$ and $\varepsilon \in (0, 1]$ we have

$$d_{GH}(\mathbb{P}(\varepsilon b), \mathbb{P}_{d_b}^m(\varepsilon, x_b)) = O(\varepsilon)$$

The function $O(\varepsilon)$ may change with b . We used the notations

$$\mathbb{P}(b) = [\bar{B}(x, 1), d_b, x_b] \quad \text{and} \quad \mathbb{P}_{d_b}^m(\varepsilon, x) = \left[\bar{B}(x, 1), \frac{1}{\varepsilon}d_b, x_b \right]$$

2.5 Curvdimension and curvature

In the case of a riemannian manifold (X, g) , with smooth enough (typically \mathcal{C}^1) metric g , the tangent metric spaces exist for any point of the manifold. Ideed, the tangent metric space $[T_x X, d^x, x]$ is the isometry class of a small neighbourhood of the origin of the tangent space (in differential geometric sense) $T_x X$, with d^x being the euclidean distance induced by the norm given by g_x . Moreover, we have the following description of the sectional curvature.

Proposition 2.22 *Let (X, d) be a \mathcal{C}^4 smooth riemannian manifold with d the length distance induced by the riemannian metric g . Suppose that for a point $x \in X$ the sectional curvature is bounded in the sense that for any linearly independent $u, v \in T_x X$ we have $|K_x(u, v)| \leq C$. Then for any sufficiently small $\varepsilon > 0$ we have*

$$\frac{1}{\varepsilon^2} d_{GH}(\mathbb{P}^m(\varepsilon, x), [T_x X, d^x, x]) \leq \frac{1}{3} C + \mathcal{O}(\varepsilon) \quad (8)$$

Proof. This is well known, in another form. Indeed, for small enough ε , consider the geodesic exponential map which associates to any $u \in W \subset T_x X$ (in a neighbourhood W of the origin which is independent of ε) the point $\exp_x \varepsilon u$. Define now the distance

$$d_\varepsilon^x(u, v) = \frac{1}{\varepsilon} d(\exp_x \varepsilon u, \exp_x \varepsilon v)$$

We can choose the neighbourhood W to be the unit ball with respect to the distance d^x in order to get the following estimate:

$$\sup \{ |d_\varepsilon^x(u, v) - d^x(u, v)| : u, v \in W \} \geq d_{GH}(\mathbb{P}^m(\varepsilon, x), [W, d^x, 0])$$

where $d^x(u, v) = \|u - v\|_x$. In the given regularity settings, we shall use the following expansion of d_ε^x : if u, v are linearly independent then

$$d_\varepsilon^x(u, v) = d^x(u, v) - \frac{1}{6} \varepsilon^2 K_x(u, v) \frac{\|u\|_x^2 \|v\|_x^2 - \langle u, v \rangle_x^2}{d^x(u, v)} + \varepsilon^2 \mathcal{O}(\varepsilon) \quad (9)$$

where K is the sectional curvature of the metric g . (If u, v are linearly dependent then $d_\varepsilon^x(u, v) = d^x(u, v)$.) From here we easily obtain that

$$\frac{1}{\varepsilon^2} d_{GH}(\mathbb{P}^m(\varepsilon, x), [T_x X, d^x, x]) \leq \frac{1}{3} \sup \{ |K_x(u, v)| : u, v \in W \text{ lin. indep.} \} + \mathcal{O}(\varepsilon)$$

which ends the proof. \square

This proposition makes us define the "curvdimension" and "curvature" of an (abstract) metric profile.

Definition 2.23 *Let \mathbb{P} be an abstract metric profile. The curvdimension of this abstract metric profile is*

$$\text{curvdim } \mathbb{P} = \sup \left\{ \alpha > 0 : \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^\alpha} d_{GH}(\mathbb{P}(\varepsilon), \mathbb{P}(0)) = 0 \right\} \quad (10)$$

and, in the case that the curvdimension equals $\beta > 0$ then the β -curvature of \mathbb{P} is the number $M > 0$ such that

$$\lim_{\varepsilon \rightarrow 0} \log_\varepsilon \left(\frac{1}{M} d_{GH}(\mathbb{P}(\varepsilon), \mathbb{P}(0)) \right) = \beta \quad (11)$$

In case \mathbb{P} is the metric profile of a point $x \in X$ in a metric space (X, d) then the curvdimension at x and curvature at x are the curvdimension, respectively curvature, of the metric profile of x .

It follows that non-flat riemannian (smooth enough) spaces have curvdimension 2. Also, any metric cone has curvdimension equal to 0 (meaning "all metric cones are flat objects"). In particular, finite dimensional normed vector spaces are flat (as they should be).

Question. What is the curvdimension of a sub-riemannian space? Carnot groups endowed with left invariant Carnot-Carathéodory distances (or any left-invariant distance coming from a norm on the Carnot group seen as a conical group) have curvdimension equal to zero, i.e. they are "flat". In the non-flat case, that is when the metric profiles are not constant, what is then the curvdimension at a generic point of a sub-riemannian space? In the paper [11], theorem 10.1, then in section 8, [13], we presented evidence for the fact that metric contact 3 dimensional manifolds have curvdimension smaller than 2. Is it, in this case, equal to 1?

3 Groups with dilations

We shall see that for a dilation structure (or "dilatation structure", or "metric space with dilations") the metric tangent spaces have the structure of a normed local group with dilations. The notion has been introduced in published [10], [4]; it appears first time in [12], which we follow here.

We shall work with local groups and local actions instead of the usual global ones. We shall use "uniform local group" for a local group endowed with its canonical uniform structure.

Let Γ be a topological commutative group, endowed with a continuous morphism $|\cdot|: \Gamma \rightarrow (0, +\infty)$. For example Γ could be $(0, +\infty)$ with the operation of multiplication of positive real numbers and the said morphism could be the identity. Or Γ could be the set of complex numbers different from 0, with the operation of multiplication of complex numbers and morphism taken to be the modulus function. Also, Γ could be the set of integers with the operation of addition and the morphism could be the exponential function. Many other possibilities exist (like a product between a finite commutative group with one of the examples given before).

It is useful further to just think that Γ is like in the first example, because in these notes we are not going to use the structure of Γ in order to put more geometrical objects on the metric space (like we do, for example, in the paper [8]).

The elements of Γ will be denoted with small greek letters, like ε, μ, \dots . By covention, whenever we write " $\varepsilon \rightarrow 0$ ", we really mean " $|\varepsilon| \rightarrow 0$ ". Also " $\mathcal{O}(\varepsilon)$ " means " $\mathcal{O}(|\varepsilon|)$ ", and so on.

Definition 3.1 (G, δ) is a local group with dilations if G is a local group and $\delta: \Gamma \rightarrow \mathcal{C}(G, G)$ is a local action of Γ on G , such that

H0. the following limit $\lim_{\varepsilon \rightarrow 0} \delta_\varepsilon x = e$ is uniform with respect to x in a compact neighbourhood of the identity element e .

H1. the limit $\beta(x, y) = \lim_{\varepsilon \rightarrow 0} \delta_\varepsilon^{-1}((\delta_\varepsilon x)(\delta_\varepsilon y))$ is is uniform with respect to (x, y) in a compact neighbourhood of (e, e) .

H2. the limit $\lim_{\varepsilon \rightarrow 0} \delta_\varepsilon^{-1}((\delta_\varepsilon x)^{-1}) = x^{-1}$ is uniform with respect to x in a compact neighbourhood of the identity element e .

Definition 3.2 A normed local group with dilations $(G, \|\cdot\|, \delta)$ is a local group with dilations (G, δ) endowed with a continuous norm, $\|\cdot\|: G \rightarrow [0, +\infty)$ which satisfies:

(a) there is a function $\|\cdot\|: U \subset G \rightarrow [0, +\infty)$ defined on a neighbourhood U of e , such that the limit $\lim_{\varepsilon \rightarrow 0} \frac{1}{|\varepsilon|} \|\delta_\varepsilon x\| = \|x\|^N$ is uniform with respect to x in compact set,

(b) if $\|x\|^N = 0$ then $x = e$.

In a normed local group with dilations we consider the left invariant (locally defined) distance given by

$$d(x, y) = \|x^{-1}y\| \quad . \quad (12)$$

and dilations based in any point $x \in G$ by

$$\delta_\varepsilon^x u = x \delta_\varepsilon(x^{-1}u). \quad (13)$$

3.1 Conical groups

Definition 3.3 A normed conical group $(N, \|\cdot\|, \delta)$ is a normed group with dilations such that for any $\varepsilon \in \Gamma$: (a) the dilation δ_ε is a group morphism and (b) the norm is homogeneous, that is $\|\delta_\varepsilon x\| = |\varepsilon| \|x\|$.

Then, a conical group appears as the infinitesimal version of a group with dilations ([4] proposition 2). For the proof see the more general theorem 7.3.

Proposition 3.4 Let $(G, \|\cdot\|, \delta)$ be a normed local group with dilations. Then $(G, \|\cdot\|^N, \delta)$ is a local normed conical group, with operation β , dilations δ and homogeneous norm $\|\cdot\|^N$.

3.2 Carnot groups

Carnot groups appear in sub-riemannian geometry as models of tangent spaces, [3], [25], [29]. In particular such groups can be endowed with a structure of a sub-riemannian manifold. Here we are interested in the fact that they are particular examples of conical groups.

Definition 3.5 A Carnot group is a pair (N, V_1) formed by a real connected simply connected Lie group N , with a distinguished subspace V_1 of the Lie algebra $Lie(N)$, such that

$$Lie(N) = \sum_{i=1}^m V_i, \quad V_{i+1} = [V_1, V_i]$$

The number m is called the step of the group. The number $Q = \sum_{i=1}^m i \dim V_i$ is called the homogeneous dimension of the group.

Because the group N is nilpotent and simply connected, it follows that the (Lie group) exponential mapping is a diffeomorphism. It is customary to identify then the group with the algebra. We obtain a set N equal to some \mathbb{R}^n , endowed with a Lie algebra structure (that is a real vector space and a Lie bracket) and a Lie group structure (that is a Lie group operation, denoted multiplicatively, defined for any pair of elements of N , with the 0 element of the vector space N as neutral element for the group operation).

The Baker-Campbell-Hausdorff formula connects the Lie bracket and the group operation. Indeed, the algebra being nilpotent, it follows that the group operation is polynomial, because the Baker-Campbell-Hausdorff formula contains only a finite number of terms. Thus, the group operation is expressed as a function of the Lie bracket operation. Moreover, Lie algebra endomorphisms are group endomorphisms and the converse is also true.

The most simple case is when the Lie bracket is a constant function equal to 0 and $V_1 = N = \mathbb{R}^n$. In this case the group operation is the vector space addition (even if defined multiplicatively). The step of this group is equal to 1 and the homogeneous dimension equals n .

Let us take $\Gamma = (0, +\infty)$, $|\cdot|: \Gamma \rightarrow (0, \infty)$ the density function and let us define for any $\varepsilon > 0$, the dilation:

$$x = \sum_{i=1}^m x_i \mapsto \delta_\varepsilon x = \sum_{i=1}^m \varepsilon^i x_i$$

Any such dilation is a group morphism and a Lie algebra morphism.

Let us choose an euclidean norm $\|\cdot\|$ on V_1 . We shall endow the group N with a norm coming from a Carnot-Carathéodory distance (general definition in section 9.1). Remark that for any $x \in V_1$ and any $\varepsilon > 0$ we have $\|\delta_\varepsilon x\| = \varepsilon \|x\|$.

Indeed, by definition the space V_1 generates N (as a Lie algebra), therefore any element $x \in N$ can be written as a product of elements from V_1 . A controlled way to do this is described in the following slight reformulation of Lemma 1.40, Folland, Stein [21]).

Lemma 3.6 *Let N be a Carnot group and X_1, \dots, X_p an orthonormal basis for V_1 . Then there is a natural number M and a function $g: \{1, \dots, M\} \rightarrow \{1, \dots, p\}$ such that any $x \in N$ can be written as:*

$$x = \prod_{i=1}^M \exp(t_i X_{g(i)}) \tag{14}$$

Moreover, if x is sufficiently close (in Euclidean norm) to 0 then each t_i can be chosen such that $|t_i| \leq C \|x\|^{1/m}$

From these data we may construct a norm on the Carnot group N , by the intermediary of a Carnot-Carathéodory (CC for short) distance. Here we give an algebraic definition of this distance.

The (Carnot-Carathéodory) norm on the Carnot group is defined as

$$\|x\| = \inf \left\{ \sum_{i \in I} \|x_i\| : \text{all finite sets } I \text{ and all decompositions } x = \prod_{i \in I} x_i \text{ where all } x_i \in V_1 \right\}$$

The CC norm is then finite (by lemma 3.6) for any two $x \in N$ and it is also continuous. All in all $(N, \|\cdot\|, \delta)$ is a (global) normed conical group.

3.3 Contractible groups

Definition 3.7 *A contractible group is a pair (G, α) , where G is a topological group with neutral element denoted by e , and $\alpha \in \text{Aut}(G)$ is an automorphism of G such that:*

- α is continuous, with continuous inverse,
- for any $x \in G$ we have the limit $\lim_{n \rightarrow \infty} \alpha^n(x) = e$.

For a contractible group (G, α) , the automorphism α is compactly contractive (Lemma 1.4 (iv) [30]), that is: for each compact set $K \subset G$ and open set $U \subset G$, with $e \in U$, there is $N(K, U) \in \mathbb{N}$ such that for any $x \in K$ and $n \in \mathbb{N}$, $n \geq N(K, U)$, we have $\alpha^n(x) \in U$.

If G is locally compact then α compactly contractive is equivalent with: each identity neighbourhood of G contains an α -invariant neighbourhood. Further on we shall assume without mentioning that all groups are locally compact.

Any conical group is a contractible group. Indeed, it suffices to associate to a conical group (G, δ) the contractible group (G, δ_ε) , for a fixed $\varepsilon \in \Gamma$ with $\nu(\varepsilon) < 1$.

Conversely, to any contractible group (G, α) we may associate the conical group (G, δ) , with $\Gamma = \left\{ \frac{1}{2^n} : n \in \mathbb{N} \right\}$ and for any $n \in \mathbb{N}$ and $x \in G$

$$\delta_{\frac{1}{2^n}} x = \alpha^n(x) \quad .$$

(Finally, a local conical group has only locally the structure of a contractible group.)

The structure of contractible groups is known to some detail, due to results by Siebert [30], Wang [35], Glöckner and Willis [24], Glöckner [23] and others (see references in the mentioned papers).

Related to contractible groups, here is the definition of a contractive automorphism group [30], definition 5.1.

Definition 3.8 *Let G be a locally compact group. An automorphism group on G is a family $T = (\tau_t)_{t>0}$ in $\text{Aut}(G)$, such that $\tau_t \tau_s = \tau_{ts}$ for all $t, s > 0$.*

The contraction group of T is defined by

$$C(T) = \left\{ x \in G : \lim_{t \rightarrow 0} \tau_t(x) = e \right\} \quad .$$

The automorphism group T is contractive if $C(T) = G$.

It is obvious that a contractive automorphism group T induces on G a structure of conical group. Conversely, any conical group with $\Gamma = (0, +\infty)$ has an associated contractive automorphism group (the group of dilations based at the neutral element).

Siebert, proposition 5.4 [30], gives a very useful description of a class of contractible groups.

Proposition 3.9 *For a locally compact group G the following assertions are equivalent:*

- (i) *G admits a contractive automorphism group;*
- (ii) *G is a simply connected Lie group whose Lie algebra admits a positive graduation.*

These groups are almost Carnot groups. Indeed, what is missing is the fact that the first element of the graduation generates the Lie algebra.

4 Dilation structures

In this paper I use the denomination "dilation structure", or "metric space with dilations", compared with older papers, where the name "dilatation structure" was used.

We shall use here a slightly particular version of dilation structures. For the general definition of a dilation structure see [4] (the general definition applies for dilation structures over ultrametric spaces as well).

4.1 Normed groupoids with dilations

Notions of convergence. We need a topology on a normed groupoid (G, d) , induced by the norm.

Definition 4.1 *A net of arrows (a_ε) simply converges to the arrow $a \in G$ (we write $a_\varepsilon \rightarrow a$) if:*

- (i) *for any $\varepsilon \in I$ there are elements $g_\varepsilon, h_\varepsilon \in G$ such that $h_\varepsilon a_\varepsilon g_\varepsilon = a$,*

(ii) we have $\lim_{\varepsilon \in I} d(g_\varepsilon) = 0$ and $\lim_{\varepsilon \in I} d(h_\varepsilon) = 0$.

A net of arrows (a_ε) left-converges to the arrow $a \in G$ (we write $a_\varepsilon \xrightarrow{L} a$) if for all $i \in I$ we have $(a_\varepsilon^{-1}, a) \in G^{(2)}$ and moreover $\lim_{\varepsilon \in I} d(a_\varepsilon^{-1}a) = 0$.

A net of arrows (a_ε) right-converges to the arrow $a \in G$ (we write $a_\varepsilon \xrightarrow{R} a$) if for all $i \in I$ we have $(a_\varepsilon, a^{-1}) \in G^{(2)}$ and moreover $\lim_{\varepsilon \in I} d(a_\varepsilon a^{-1}) = 0$.

It is clear that if $a_\varepsilon \xrightarrow{L} a$ or $a_\varepsilon \xrightarrow{R} a$ then $a_\varepsilon \rightarrow a$.

Right-convergence of a_ε to a is just convergence of a_ε to a in the distance $d_{\alpha(a)}$, that is $\lim_{\varepsilon \in I} d_{\alpha(a)}(a_\varepsilon, a) = 0$.

Left-convergence of a_ε to a is just convergence of a_ε^{-1} to a^{-1} in the distance $d_{\omega(a)}$, that is $\lim_{\varepsilon \in I} d_{\omega(a)}(a_\varepsilon^{-1}, a^{-1}) = 0$.

Proposition 4.2 *Let (G, d) be a normed groupoid.*

(i) If $a_\varepsilon \xrightarrow{L} a$ and $a_\varepsilon \xrightarrow{L} b$ then $a = b$. If $a_\varepsilon \xrightarrow{R} a$ and $a_\varepsilon \xrightarrow{R} b$ then $a = b$.

(ii) The following are equivalent:

1. G is a Hausdorff topological groupoid with respect to the topology induced by the simple convergence,
2. d is a separable norm,
3. for any net (a_ε) , if $a_\varepsilon \rightarrow a$ and $a_\varepsilon \rightarrow b$ then $a = b$.
4. for any net (a_ε) , if $a_\varepsilon \xrightarrow{R} a$ and $a_\varepsilon \xrightarrow{L} b$ then $a = b$.

Proof. (i) We prove only the first part of the conclusion. We can write $b^{-1}a = b^{-1}a_\varepsilon a_\varepsilon^{-1}a$, therefore

$$d(b^{-1}a) \leq d(b^{-1}a_\varepsilon) + d(a_\varepsilon^{-1}a)$$

The right hand side of this inequality is arbitrarily small, so $d(b^{-1}a) = 0$, which implies $a = b$.

(ii) Remark that the structure maps are continuous with respect to the topology induced by the simple convergence. We need only to prove the uniqueness of limits.

3. \Rightarrow 4. is trivial. In order to prove that 4. \Rightarrow 3., consider an arbitrary net (a_ε) such that $a_\varepsilon \rightarrow a$ and $a_\varepsilon \rightarrow b$. This means that there exist nets $(g_\varepsilon), (g'_\varepsilon), (h_\varepsilon), (h'_\varepsilon)$ such that $h_\varepsilon a_\varepsilon g_\varepsilon = a$, $h'_\varepsilon a_\varepsilon g'_\varepsilon = b$ and $\lim_{i \in I} (d(g_\varepsilon) + d(g'_\varepsilon) + d(h_\varepsilon) + d(h'_\varepsilon)) = 0$. Let $g''_\varepsilon = g_\varepsilon^{-1}g'_\varepsilon$ and $h''_\varepsilon = h'_\varepsilon h_\varepsilon^{-1}$. We have then $b = h''_\varepsilon a g''_\varepsilon$ and $\lim_{i \in I} (d(g''_\varepsilon) + d(h''_\varepsilon)) = 0$. Then $h''_\varepsilon a \xrightarrow{L} b$ and $h''_\varepsilon a \xrightarrow{R} a$. We deduce that $a = b$.

1. \Leftrightarrow 3. is trivial. So is 3. \Rightarrow 2. We finish the proof by showing that 2. \Rightarrow 3. By a reasoning made previously, it is enough to prove that: if $b = h_\varepsilon a g_\varepsilon$ and $\lim_{i \in I} (d(g_\varepsilon) + d(h_\varepsilon)) = 0$ then $a = b$. Because d is separable it follows that $\alpha(a) = \alpha(b)$ and $\omega(a) = \omega(b)$. We have then $a^{-1}b = a^{-1}h_\varepsilon a g_\varepsilon$, therefore

$$d(a^{-1}b) \leq d(a^{-1}h_\varepsilon a) + d(g_\varepsilon)$$

The norm d induces a left invariant distance on the vertex group of all arrows g such that $\alpha(g) = \omega(g) = \alpha(a)$. This distance is obviously continuous with respect to the simple convergence in the group. The net $a^{-1}h_\varepsilon a$ simply converges to $\alpha(a)$ by the continuity of the multiplication (indeed, h_ε

simply converges to $\alpha(a)$). Therefore $\lim_{i \in I} d(a^{-1}h_\varepsilon a) = 0$. It follows that $d(a^{-1}b)$ is arbitrarily small, therefore $a = b$. \square

By adapting the definition of a normed group with dilations to a normed groupoid with dilations, we get the following structure.

Definition 4.3 *A normed groupoid (G, d, δ) with dilations is a separated normed groupoid (G, d) endowed with a map assigning to any $\varepsilon \in \Gamma$ a transformation $\delta_\varepsilon : \text{dom}(\varepsilon) \rightarrow \text{im}(\varepsilon)$ which satisfies the following:*

- A1. *For any $\varepsilon \in \Gamma$ $\alpha\delta_\varepsilon = \alpha$. Moreover $\varepsilon \in \Gamma \mapsto \delta_\varepsilon$ is an action of Γ on G , that is for any $\varepsilon, \mu \in \Gamma$ we have $\delta_\varepsilon\delta_\mu = \delta_{\varepsilon\mu}$, $(\delta_\varepsilon)^{-1} = \delta_{\varepsilon^{-1}}$ and $\delta_e = \text{id}$.*
- A2. *For any $x \in \text{Ob}(G)$ and any $\varepsilon \in \Gamma$ we have $\delta_\varepsilon(x) = x$. Moreover the transformation δ_ε contracts $\text{dom}(\varepsilon)$ to $X = \text{Ob}(G)$ uniformly on bounded sets, which means that the net $d\delta_\varepsilon$ converges to the constant function 0, uniformly on bounded sets.*
- A3. *There is a function $\bar{d} : U \rightarrow \mathbb{R}$ which is the limit*

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{|\varepsilon|} d\delta_\varepsilon(g) = \bar{d}(g)$$

uniformly on bounded sets in the sense of definition 4.1. Moreover, if $\bar{d}(g) = 0$ then $g \in \text{Ob}(G)$.

- A4. *the net dif_ε converges uniformly on bounded sets to a function $\bar{\text{dif}}$.*

The domains and codomains of a dilation of (G, d) satisfy the following Axiom A0:

- (i) *for any $\varepsilon \in \Gamma$ $\text{Ob}(G) = X \subset \text{dom}(\varepsilon)$ and $\text{dom}(\varepsilon) = \text{dom}(\varepsilon)^{-1}$,*
- (ii) *for any bounded set $K \subset \text{Ob}(G)$ there are $1 < A < B$ such that for any $\varepsilon \in \Gamma$, $|\varepsilon| \leq 1$:*

$$\begin{aligned} d^{-1}(|\varepsilon|) \cap \alpha^{-1}(K) &\subset \delta_\varepsilon(d^{-1}(A) \cap \alpha^{-1}(K)) \subset \text{dom}(\varepsilon^{-1}) \cap \alpha^{-1}(K) \subset \\ &\subset \delta_\varepsilon(d^{-1}(B) \cap \alpha^{-1}(K)) \subset \delta_\varepsilon(\text{dom}(\varepsilon) \cap \alpha^{-1}(K)) \end{aligned} \quad (15)$$

- (iii) *for any bounded set $K \subset \text{Ob}(G)$ there are $R > 0$ and $\varepsilon_0 \in (0, 1]$ such that for any $\varepsilon \in \Gamma$, $|\varepsilon| \leq \varepsilon_0$ and any $g, h \in d^{-1}(R) \cap \alpha^{-1}(K)$ we have:*

$$\text{dif}(\delta_\varepsilon g, \delta_\varepsilon h) \in \text{dom}(\varepsilon^{-1}) \quad (16)$$

4.2 Dilation structures, definition

By proposition 2.5, any metric space (X, d) may be seen as the normed groupoid $(X \times X, d)$. Let us see what happens if we endow this trivial groupoid with dilations, according to definition 4.3. For any $\varepsilon \in \Gamma$ we have a dilation

$$\delta_\varepsilon : \text{dom}(\varepsilon) \subset X^2 \rightarrow \text{im}(\varepsilon) \subset X^2$$

which satisfies a number of axioms. Let us take them one by one.

A1. for trivial groupoids. For any $\varepsilon \in \Gamma$ $\alpha\delta_\varepsilon = \alpha$ is equivalent to the existence of a locally defined function δ_ε^x , for any $x \in X$, such that

$$\delta_\varepsilon(y, x) = (\delta_\varepsilon^x y, x)$$

The domain of definition of δ_ε^x is $\text{dom}(\varepsilon) \cap \{(y, x) : y \in X\}$ (similarly for the image). The fact that $\varepsilon \in \Gamma \mapsto \delta_\varepsilon$ is an action of Γ on X^2 , translates into: for any $\varepsilon, \mu \in \Gamma$ and $x \in X$ we have $\delta_\varepsilon^x \delta_\mu^x = \delta_{\varepsilon\mu}^x$, $(\delta_\varepsilon^x)^{-1} = \delta_{\varepsilon^{-1}}^x$ and $\delta_e^x = \text{id}$.

A2. for trivial groupoids. The objects of the trivial groupoid X^2 are of the form (x, x) with $x \in X$. We use what we already know from A1 to deduce that the axiom A2 says: for any $x \in X$ and any $\varepsilon \in \Gamma$ we have $\delta_\varepsilon^x x = x$. Moreover the transformation $(y, x) \mapsto (\delta_\varepsilon^x y, x)$ contracts the domain $\text{dom}(\varepsilon)$ to $\{(x, x) : x \in X\}$, uniformly on sets $A \subset X^2$ which are "bounded" in the sense: there is a $M > 0$ such that for any $(x, y) \in A$ we have $d(x, y) \leq M$. This means that the net of functions $(x, y) \mapsto d(\delta_\varepsilon^x y, x)$ converges to the constant function 0, uniformly with respect to $(x, y) \in A$, where A is bounded in the sense explained before.

A3. for trivial groupoids. A simple computation shows that a pair $(g, h) \in G \times_\alpha G$ has the form $g = (u, x)$, $h = (v, x)$. Then

$$\text{dif}(\delta_\varepsilon g, \delta_\varepsilon h) = \left(\delta_{\varepsilon^{-1}}^{\delta_\varepsilon^x v} \delta_\varepsilon^x u \right)$$

The axiom A3 says that for any $x \in X$ there is a function d^x , locally defined on pair of points $(u, v) \in X \times X$, such that

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{|\varepsilon|} d \left(\delta_{\varepsilon^{-1}}^{\delta_\varepsilon^x v} \delta_\varepsilon^x u \right) = d_x(u, v)$$

uniformly with respect to $d(x, u)$, $d(x, v)$. Moreover, if $d_x(u, v) = 0$ then $u = v$.

A4. for trivial groupoids. Using A2 for trivial groupoids, this axiom says that the net $\delta_{\varepsilon^{-1}}^{\delta_\varepsilon^x v} \delta_\varepsilon^x u$ converges to a function $\Delta^x(v, u)$ uniformly with respect to $d(x, u)$, $d(x, v)$.

The axiom A0 can also be detailed. All in all we see that trivial normed groupoids with dilations correspond to strong dilation structures. defined next.

Definition 4.4 Let (X, d) be a complete metric space such that for any $x \in X$ the closed ball $\bar{B}(x, 3)$ is compact. A dilation structure (X, d, δ) over (X, d) is the assignment to any $x \in X$ and $\varepsilon \in (0, +\infty)$ of a invertible homeomorphism, defined as: if $\varepsilon \in (0, 1]$ then $\delta_\varepsilon^x : U(x) \rightarrow V_\varepsilon(x)$, else $\delta_\varepsilon^x : W_\varepsilon(x) \rightarrow U(x)$, such that the following axioms are satisfied:

A0. For any $x \in X$ the sets $U(x), V_\varepsilon(x), W_\varepsilon(x)$ are open neighbourhoods of x . There are numbers $1 < A < B$ such that for any $x \in X$ and any $\varepsilon \in (0, 1)$ we have the following string of inclusions:

$$B_d(x, \varepsilon) \subset \delta_\varepsilon^x B_d(x, A) \subset V_\varepsilon(x) \subset W_{\varepsilon^{-1}}(x) \subset \delta_\varepsilon^x B_d(x, B)$$

Moreover for any compact set $K \subset X$ there are $R = R(K) > 0$ and $\varepsilon_0 = \varepsilon(K) \in (0, 1)$ such that for all $u, v \in \bar{B}_d(x, R)$ and all $\varepsilon \in (0, \varepsilon_0)$, we have

$$\delta_\varepsilon^x v \in W_{\varepsilon^{-1}}(\delta_\varepsilon^x u) .$$

A1. We have $\delta_\varepsilon^x x = x$ for any point x . We also have $\delta_1^x = \text{id}$ for any $x \in X$. Let us define the topological space

$$\text{dom } \delta = \{(\varepsilon, x, y) \in (0, +\infty) \times X \times X : \text{if } \varepsilon \leq 1 \text{ then } y \in U(x) \},$$

else $y \in W_\varepsilon(x)$

with the topology inherited from $(0, +\infty) \times X \times X$ endowed with the product topology. Consider also $Cl(dom \delta)$, the closure of $dom \delta$ in $[0, +\infty) \times X \times X$. The function $\delta : dom \delta \rightarrow X$ defined by $\delta(\varepsilon, x, y) = \delta_\varepsilon^x y$ is continuous. Moreover, it can be continuously extended to the set $Cl(dom \delta)$ and we have

$$\lim_{\varepsilon \rightarrow 0} \delta_\varepsilon^x y = x$$

A2. For any $x \in X$, $\varepsilon, \mu \in (0, +\infty)$ and $u \in U(x)$ we have the equality:

$$\delta_\varepsilon^x \delta_\mu^x u = \delta_{\varepsilon\mu}^x u$$

whenever one of the sides are well defined.

A3. For any x there is a distance function $(u, v) \mapsto d^x(u, v)$, defined for any u, v in the closed ball (in distance d) $\bar{B}(x, A)$, such that

$$\lim_{\varepsilon \rightarrow 0} \sup \left\{ \left| \frac{1}{\varepsilon} d(\delta_\varepsilon^x u, \delta_\varepsilon^x v) - d^x(u, v) \right| : u, v \in \bar{B}_d(x, A) \right\} = 0$$

uniformly with respect to x in compact set.

The dilation structure is strong if it satisfies the following supplementary condition:

A4. Let us define $\Delta_\varepsilon^x(u, v) = \delta_{\varepsilon^{-1}}^{\delta_\varepsilon^x u} \delta_\varepsilon^x v$. Then we have the limit

$$\lim_{\varepsilon \rightarrow 0} \Delta_\varepsilon^x(u, v) = \Delta^x(u, v)$$

uniformly with respect to x, u, v in compact set.

We shall use many times from now the words "sufficiently close". This deserves a definition.

Definition 4.5 Let (X, d, δ) be a strong dilation structure. We say that a property

$$\mathcal{P}(x_1, x_2, x_3, \dots)$$

is true for x_1, x_2, x_3, \dots **sufficiently close** if for any compact, non empty set $K \subset X$, there is a positive constant $C(K) > 0$ such that $\mathcal{P}(x_1, x_2, x_3, \dots)$ is true for any $x_1, x_2, x_3, \dots \in K$ with $d(x_i, x_j) \leq C(K)$.

5 Examples of dilation structures

5.1 Snowflakes, nonstandard dilations in the plane

Snowflake construction. This is an adaptation of a standard construction for metric spaces with dilations: if (X, d, δ) is a dilation structure then $(X, d_a, \delta(a))$ is also a dilation structure, for any $a \in (0, 1]$, where

$$d_a(x, y) = (d(x, y))^a, \quad \delta(a)_\varepsilon^x = \delta_{\varepsilon^{\frac{1}{a}}}^x.$$

In particular, if $X = \mathbb{R}^n$ then for any $a \in (0, 1]$ we may take the distance and dilations

$$d_a(x, y) = \|x - y\|^a, \quad \delta_\varepsilon^x y = x + \varepsilon^{\frac{1}{a}}(y - x).$$

Nonstandard dilations. In the plane $X = \mathbb{R}^2$, endowed with the euclidean distance, we may consider another one-parameter group of linear transformations instead of the familiar homotheties. Indeed, for any complex number $z = 1 + i\theta$ let us define the dilations

$$\delta_\varepsilon x = \varepsilon^z x .$$

Then $(X, \delta, +, d)$ is a conical group, therefore the dilations

$$\delta_\varepsilon^x y = x + \delta_\varepsilon(y - x)$$

together with the euclidean distance, form a dilation structure.

Two such dilation structures, constructed respectively by using the numbers $1 + i\theta$ and $1 + i\theta'$, are equivalent (see the section 7.4 for the definition of equivalent dilation structures) if and only if $\theta = \theta'$.

Such dilation structures give examples of metric spaces with dilations which don't have the Radon-Nikodym property, see section 8.1.

5.2 Normed groups with dilations

The following result is theorem 15 [4].

Theorem 5.1 *Let $(G, \delta, \|\cdot\|)$ be a locally compact normed local group with dilations. Then (G, d, δ) is a dilation structure, where δ are the dilations defined by (13) and the distance d is induced by the norm as in (12).*

Proof. The axiom A0 is straightforward from definition 3.1, axiom H0, and because the dilation structure is left invariant, in the sense that the transport by left translations in G preserves the dilations δ . We also trivially have axioms A1 and A2 satisfied.

For the axiom A3 remark that $d(\delta_\varepsilon^x u, \delta_\varepsilon^x v) = d(x\delta_\varepsilon(x^{-1}u), x\delta_\varepsilon(x^{-1}v)) = d(\delta_\varepsilon(x^{-1}u), \delta_\varepsilon(x^{-1}v))$. Let us denote $U = x^{-1}u$, $V = x^{-1}v$ and for $\varepsilon > 0$ let

$$\beta_\varepsilon(u, v) = \delta_\varepsilon^{-1}((\delta_\varepsilon u)(\delta_\varepsilon v)).$$

We have then:

$$\frac{1}{\varepsilon} d(\delta_\varepsilon^x u, \delta_\varepsilon^x v) = \frac{1}{\varepsilon} \|\delta_\varepsilon \beta_\varepsilon(\delta_\varepsilon^{-1}((\delta_\varepsilon V)^{-1}), U)\| .$$

Define the function

$$d^x(u, v) = \|\beta(V^{-1}, U)\|^N .$$

From definition 3.1 axioms H1, H2, and from definition 3.2 (d), we obtain that axiom A3 is satisfied.

For the axiom A4 we have to compute:

$$\begin{aligned} \Delta^x(u, v) &= \delta_{\varepsilon^{-1}}^{\delta_\varepsilon^x u} \delta_\varepsilon^x v = (\delta_\varepsilon^x u)(\delta_\varepsilon)^{-1} \left((\delta_\varepsilon^x u)^{-1} (\delta_\varepsilon^x v) \right) = \\ &= (x\delta_\varepsilon U) \beta_\varepsilon(\delta_\varepsilon^{-1}((\delta_\varepsilon V)^{-1}), U) \rightarrow x\beta(V^{-1}, U) \end{aligned}$$

as $\varepsilon \rightarrow 0$. Therefore axiom A4 is satisfied. \square

5.3 Riemannian manifolds

The following interesting quotation from Gromov book [26], pages 85-86, motivates some of the ideas underlying dilation structures, especially in the very particular case of a riemannian manifold:

“3.15. Proposition: *Let (V, g) be a Riemannian manifold with g continuous. For each $v \in V$ the spaces $(V, \lambda d, v)$ Lipschitz converge as $\lambda \rightarrow \infty$ to the tangent space $(T_v V, 0)$ with its Euclidean metric g_v .*

Proof₊: *Start with a C^1 map $(\mathbb{R}^n, 0) \rightarrow (V, v)$ whose differential is isometric at 0. The λ -scalings of this provide almost isometries between large balls in \mathbb{R}^n and those in λV for $\lambda \rightarrow \infty$. **Remark:** *In fact we can define Riemannian manifolds as locally compact path metric spaces that satisfy the conclusion of Proposition 3.15.*“*

The problem of domains and codomains left aside, any chart of a Riemannian manifold induces locally a dilation structure on the manifold. Indeed, take (M, d) to be a n -dimensional Riemannian manifold with d the distance on M induced by the Riemannian structure. Consider a diffeomorphism ϕ of an open set $U \subset M$ onto $V \subset \mathbb{R}^n$ and transport the dilations from V to U (equivalently, transport the distance d from U to V). There is only one thing to check in order to see that we got a dilation structure: the axiom A3, expressing the compatibility of the distance d with the dilations. But this is just a metric way to express the distance on the tangent space of M at x as a limit of rescaled distances (see Gromov Proposition 3.15, [26], p. 85-86). Denoting by g_x the metric tensor at $x \in U$, we have:

$$\begin{aligned} & [d^x(u, v)]^2 = \\ & = g_x \left(\frac{d}{d\varepsilon}|_{\varepsilon=0} \phi^{-1}(\phi(x) + \varepsilon(\phi(u) - \phi(x))), \frac{d}{d\varepsilon}|_{\varepsilon=0} \phi^{-1}(\phi(x) + \varepsilon(\phi(v) - \phi(x))) \right) \end{aligned}$$

A different example of a dilation structure on a riemannian manifold comes from the setting of proposition 2.22, section 2.5.

Let M be a smooth enough riemannian manifold and \exp be the geodesic exponential. To any point $x \in M$ and any vector $v \in T_x M$ the point $\exp_x(v) \in M$ is located on the geodesic passing thru x and tangent to v ; if we parameterize this geodesic with respect to length, such that the tangent at x is parallel and has the same direction as v , then $\exp_x(v) \in M$ has the coordinate equal with the length of v with respect to the norm on $T_x M$. We define implicitly the dilation based at x , of coefficient $\varepsilon > 0$ by the relation:

$$\delta_\varepsilon^x \exp_x(u) = \exp_x(\varepsilon u) \quad .$$

Proposition 5.2 *The above example is a strong dilation structure.*

Proof. This field of dilations satisfies trivially A0, A1, A2, only A3 and A4 are left to be checked.

Proposition 2.22 provides a proof for A3. For the proof of A4 see the section 9, where normal and adapted frames are defined for sub-riemannian manifolds. In particular the same construction works for riemannian manifolds, where one can attach normal frames to geodesic coordinate systems. \square

6 Length dilation structures

Consider (X, d) a complete, locally compact metric space, and a triple (X, d, δ) which satisfies A0, A1, A2. Denote by $Lip([0, 1], X, d)$ the space of d -Lipschitz curves $c : [0, 1] \rightarrow X$. Let also l_d denote the length functional associated to the distance d .

Definition 6.1 *For any $\varepsilon \in (0, 1)$ and $x \in X$ we define the length functional at scale ε , relative to x , to be*

$$l_\varepsilon(x, c) = l_\varepsilon^x(c) = \frac{1}{\varepsilon} l_d(\delta_\varepsilon^x c)$$

The domain of definition of the functional l_ε is the space:

$$\mathcal{L}_\varepsilon(X, d, \delta) = \{(x, c) \in X \times \mathcal{C}([0, 1], X) : c : [0, 1] \in U(x), \\ \delta_\varepsilon^x c \text{ is } d\text{-Lip and } \text{Lip}(\delta_\varepsilon^x c) \leq 2l_d(\delta_\varepsilon^x c)\}$$

The last condition from the definition of $\mathcal{L}_\varepsilon(X, d, \delta)$ is a selection of parameterization of the path $c([0, 1])$. Indeed, by the reparameterization theorem, if $\delta_\varepsilon^x c : [0, 1] \rightarrow (X, d)$ is a d -Lipschitz curve of length $L = l_d(\delta_\varepsilon^x c)$ then $\delta_\varepsilon^x c([0, 1])$ can be reparameterized by length, that is there exists an increasing function $\phi : [0, L] \rightarrow [0, 1]$ such that $c' = \delta_\varepsilon^x c \circ \phi$ is a d -Lipschitz curve with $\text{Lip}(c') \leq 1$. But we can use a second affine reparameterization which sends $[0, L]$ back to $[0, 1]$ and we get a Lipschitz curve c'' with $c''([0, 1]) = c'([0, 1])$ and $\text{Lip}(c'') \leq 2l_d(c)$.

In the definition of dilation structures we use uniform convergence of distances. Here we need a notion of convergence for length functionals. This is the Gamma-convergence, a notion used many times in calculus of variations. A good reference is the book [18].

Definition 6.2 Let Z be a metric space with distance function D and $(l_\varepsilon)_{\varepsilon > 0}$ be a family of functionals $l_\varepsilon : Z_\varepsilon \subset Z \rightarrow [0, +\infty]$. Then l_ε Gamma-converges to the functional $l : Z_0 \subset Z \rightarrow [0, +\infty]$ if:

(a) (**liminf inequality**) for any function $\varepsilon \in (0, \infty) \mapsto x_\varepsilon \in Z_\varepsilon$ such that $\lim_{\varepsilon \rightarrow 0} x_\varepsilon = x_0 \in Z_0$ we have

$$l(x_0) \leq \liminf_{\varepsilon \rightarrow 0} l_\varepsilon(x_\varepsilon)$$

(b) (**existence of a recovery sequence**) For any $x_0 \in Z_0$ and for any sequence $(\varepsilon_n)_{n \in \mathbb{N}}$ such that $\lim_{n \rightarrow \infty} \varepsilon_n = 0$ there is a sequence $(x_n)_{n \in \mathbb{N}}$ with $x_n \in Z_{\varepsilon_n}$ for any $n \in \mathbb{N}$, such that

$$l(x_0) = \lim_{n \rightarrow \infty} l_{\varepsilon_n}(x_n)$$

For our needs we shall take Z to be the space $X \times \mathcal{C}([0, 1], X)$ endowed with the distance

$$D((x, c), (x', c')) = \max\{d(x, x'), \sup\{d(c(t), c'(t)) : t \in [0, 1]\}\}$$

Let $\mathcal{L}(X, d, \delta)$ be the class of all $(x, c) \in X \times \mathcal{C}([0, 1], X)$ which appear as limits $(x_n, c_n) \rightarrow (x, c)$, with $(x_n, c_n) \in \mathcal{L}_{\varepsilon_n}(X, d, \delta)$, the family $(c_n)_n$ is d -equicontinuous and $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$.

Definition 6.3 A triple (X, d, δ) is a length dilation structure if (X, d) is a complete, locally compact metric space such that A0, A1, A2, are satisfied, together with the following axioms:

A3L. there is a functional $l : \mathcal{L}(X, d, \delta) \rightarrow [0, +\infty]$ such that for any $\varepsilon_n \rightarrow 0$ as $n \rightarrow \infty$ the sequence of functionals l_{ε_n} Gamma-converges to the functional l .

A4+ Let us define $\Delta_\varepsilon^x(u, v) = \delta_{\varepsilon^{-1}}^{\delta_\varepsilon^x u} \delta_\varepsilon^x v$ and $\Sigma_\varepsilon^x(u, v) = \delta_{\varepsilon^{-1}}^x \delta_\varepsilon^{\delta_\varepsilon^x u} v$. Then we have the limits

$$\lim_{\varepsilon \rightarrow 0} \Delta_\varepsilon^x(u, v) = \Delta^x(u, v)$$

$$\lim_{\varepsilon \rightarrow 0} \Sigma_\varepsilon^x(u, v) = \Sigma^x(u, v)$$

uniformly with respect to x, u, v in compact set.

Remark 6.4 For strong dilation structures the axioms $A0 - A4$ imply $A4+$, cf. corollary 9 [4]. The transformations $\Sigma_\varepsilon^x(u, \cdot)$ have the interpretation of approximate left translations in the tangent space of (X, d) at x .

For any $\varepsilon \in (0, 1)$ and any $x \in X$ the length functional l_ε^x induces a distance on $U(x)$:

$$\dot{d}_\varepsilon^x(u, v) = \inf \{l_\varepsilon^x(c) : (x, c) \in \mathcal{L}_\varepsilon(X, d, \delta), c(0) = u, c(1) = v\}$$

In the same way the length functional l from A3L induces a distance \dot{d}^x on $U(x)$.

Gamma-convergence implies that

$$\dot{d}^x(u, v) \geq \limsup_{\varepsilon \rightarrow 0} \dot{d}_\varepsilon^x(u, v) \tag{17}$$

but We don't believe that, at this level of generality, we could have equality without supplementary hypotheses. This means that, probably, there exist length dilation structures which are not strong dilation structures.

7 Properties of dilation structures

7.1 Metric profiles associated with dilation structures

In this subsection we shall look at dilation structures from the metric point of view, by using Gromov-Hausdorff distance and metric profiles.

Let us denote by (δ, ε) the distance on

$$\bar{B}_{d^x}(x, 1) = \{y \in X : d^x(x, y) \leq 1\}$$

given by

$$(\delta, \varepsilon)(u, v) = \frac{1}{\varepsilon} d(\delta_\varepsilon^x u, \delta_\varepsilon^x v) .$$

The following theorem is a generalization of the Mitchell [27] theorem 1, concerning sub-riemannian geometry.

Theorem 7.1 *Let (X, d, δ) be a dilation structure.*

For all $u, v \in X$ such that $d(x, u) \leq 1$ and $d(x, v) \leq 1$ and all $\mu \in (0, A)$ we have:

$$d^x(u, v) = \frac{1}{\mu} d^x(\delta_\mu^x u, \delta_\mu^x v) .$$

Therefore $(U(x), d^x, x)$ is a metric cone.

Moreover, if the dilation structure is strong then the curve $\varepsilon > 0 \mapsto \mathbb{P}^x(\varepsilon) = [\bar{B}_{d^x}(x, 1), (\delta, \varepsilon), x]$ is an abstract metric profile.

Finally, we have the following limit:

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \sup \{ |d(u, v) - d^x(u, v)| : d(x, u) \leq \varepsilon, d(x, v) \leq \varepsilon \} = 0 .$$

therefore if $A4$ holds then (X, d) admits a metric tangent space in x .

Proof. For fixed $\mu \in (0, 1)$ and variable $\varepsilon \in (0, 1)$ we have:

$$\left| \frac{1}{\mu} \frac{1}{\varepsilon} d(\delta_\varepsilon^x \delta_\mu^x u, \delta_\varepsilon^x \delta_\mu^x v) - d^x(u, v) \right| = \left| \frac{1}{\varepsilon \mu} d(\delta_{\varepsilon \mu}^x u, \delta_{\varepsilon \mu}^x v) - d^x(u, v) \right| .$$

We pass pass to the limit with $\varepsilon \rightarrow 0$ and we obtain the cone property of the distance d^x .

Let us prove that \mathbb{P}^x is an abstract metric profile. For this we have to compare two pointed metric spaces, namely $((\delta^x, \varepsilon \mu), \bar{B}_{d^x}(x, 1), x)$ and $\left(\frac{1}{\mu}(\delta^x, \varepsilon), \bar{B}_{\frac{1}{\mu}(\delta^x, \varepsilon)}(x, 1), x\right)$. Let $u \in X$ such that

$$\frac{1}{\mu}(\delta^x, \varepsilon)(x, u) \leq 1 .$$

From the axioms of dilation structures and the cone property we obtain the estimate:

$$\frac{1}{\varepsilon} d^x(x, \delta_\varepsilon^x u) \leq (\mathcal{O}(\varepsilon) + 1)\mu$$

therefore $d^x(x, u) \leq (\mathcal{O}(\varepsilon) + 1)\mu$. It follows that for any $u \in \bar{B}_{\frac{1}{\mu}(\delta^x, \varepsilon)}(x, 1)$ we can choose a point $w(u) \in \bar{B}_{d^x}(x, 1)$ such that

$$\frac{1}{\mu} d^x(u, \delta_\mu^x w(u)) = \mathcal{O}(\varepsilon) .$$

Then, by using twice A3, we obtain

$$\begin{aligned} & \left| \frac{1}{\mu}(\delta^x, \varepsilon)(u_1, u_2) - (\delta^x, \varepsilon \mu)(w(u_1), w(u_2)) \right| = \\ & = \left| \frac{1}{\varepsilon \mu} d(\delta_\varepsilon^x u_1, \delta_\varepsilon^x u_2) - \frac{1}{\varepsilon \mu} d(\delta_\varepsilon^x \delta_\mu^x w(u_1), \delta_\varepsilon^x \delta_\mu^x w(u_2)) \right| \leq \\ & \leq \frac{1}{\mu} \mathcal{O}(\varepsilon) + \frac{1}{\mu} \left| d^x(u_1, u_2) - d^x(\delta_\mu^x w(u_1), \delta_\mu^x w(u_2)) \right| \leq \\ & \leq \frac{1}{\mu} \mathcal{O}(\varepsilon) + \frac{1}{\mu} d^x(u_1, \delta_\mu^x w(u_1)) + \frac{1}{\mu} d^x(u_1, \delta_\mu^x w(u_2)) \leq \\ & \leq \frac{1}{\mu} \mathcal{O}(\varepsilon) + \mathcal{O}(\varepsilon) . \end{aligned}$$

This shows that the property (b) of an abstract metric profile is satisfied. For the property (a) of an abstract metric profile we do the following. By A0, for $\varepsilon \in (0, 1)$ and $u, v \in \bar{B}_d(x, \varepsilon)$ there exist $U, V \in \bar{B}_d(x, A)$ such that

$$u = \delta_\varepsilon^x U, v = \delta_\varepsilon^x V.$$

By the cone property we have

$$\frac{1}{\varepsilon} \left| d(u, v) - d^x(u, v) \right| = \left| \frac{1}{\varepsilon} d(\delta_\varepsilon^x U, \delta_\varepsilon^x V) - d^x(U, V) \right| .$$

By A2 we have

$$\left| \frac{1}{\varepsilon} d(\delta_\varepsilon^x U, \delta_\varepsilon^x V) - d^x(U, V) \right| \leq \mathcal{O}(\varepsilon) . \quad \square$$

7.2 The tangent bundle of a dilation structure

The following proposition contains the main relations between the approximate difference, sum and inverse operations. In [4] I explained these relations as appearing from the equivalent formalism using binary decorated trees.

Proposition 7.2 *Let $(X, \circ_\varepsilon)_{\varepsilon \in \Gamma}$ be a Γ -irq. Then we have the relations:*

- (a) $\Delta_\varepsilon^x(u, \Sigma_\varepsilon^x(u, v)) = v$ (difference is the inverse of sum)
- (b) $\Sigma_\varepsilon^x(u, \Delta_\varepsilon^x(u, v)) = v$ (sum is the inverse of difference)
- (c) $\Delta_\varepsilon^x(u, v) = \Sigma_\varepsilon^{x \circ_\varepsilon u}(\text{inv}_\varepsilon^x u, v)$ (difference approximately equals the sum of the inverse)
- (d) $\text{inv}_\varepsilon^{x \circ u} \text{inv}_\varepsilon^x u = u$ (inverse operation is approximately an involution)
- (e) $\Sigma_\varepsilon^x(u, \Sigma_\varepsilon^{x \circ_\varepsilon u}(v, w)) = \Sigma_\varepsilon^x(\Sigma_\varepsilon^x(u, v), w)$ (approximate associativity of the sum)
- (f) $\text{inv}_\varepsilon^x u = \Delta_\varepsilon^x(u, x)$
- (g) $\Sigma_\varepsilon^x(x, u) = u$ (neutral element at right).

The next theorem is the generalization of proposition 3.4. It is the main structure theorem for the tangent bundle associated to a dilation structure, see theorems 7, 8, 10 in [4].

Theorem 7.3 *Let (X, d, δ) be a strong dilation structure. Then for any $x \in X$ $(U(x), \Sigma^x, \delta^x)$ is a conical group. Moreover, left translations of this group are d^x isometries.*

Proof. (I.) Let us define the "infinitesimal translations"

$$L_u^x(v) = \lim_{\varepsilon \rightarrow 0} \Delta_\varepsilon^x(u, v)$$

and prove that they are d^x isometries.

From theorem 7.1 we get the limit, as $\varepsilon \rightarrow 0$:

$$\sup \left\{ \frac{1}{\varepsilon} | d(u, v) - d^x(u, v) | : d(x, u) \leq \frac{3}{2}\varepsilon, d(x, v) \leq \frac{3}{2}\varepsilon \right\} \rightarrow 0 \quad (18)$$

For any $\varepsilon > 0$ sufficiently small the points $x, \delta_\varepsilon^x u, \delta_\varepsilon^x v, \delta_\varepsilon^x w$ are close one to another. Indeed, we have $d(\delta_\varepsilon^x u, \delta_\varepsilon^x v) = \varepsilon(d^x(u, v) + \mathcal{O}(\varepsilon))$. Therefore, if we choose u, v, w such that $d^x(u, v) < 1, d^x(u, w) < 1$, then there is $\eta > 0$ such that for all $\varepsilon \in (0, \eta)$ we have

$$d(\delta_\varepsilon^x u, \delta_\varepsilon^x v) \leq \frac{3}{2}\varepsilon \quad , \quad d(\delta_\varepsilon^x u, \delta_\varepsilon^x w) \leq \frac{3}{2}\varepsilon \quad .$$

We use ((18)) for the basepoint $\delta_\varepsilon^x u$ to get, as $\varepsilon \rightarrow 0$

$$\frac{1}{\varepsilon} | d(\delta_\varepsilon^x v, \delta_\varepsilon^x w) - d^{\delta_\varepsilon^x u}(\delta_\varepsilon^x v, \delta_\varepsilon^x w) | \rightarrow 0$$

From the cone property of the distance $d^{\delta_\varepsilon^x u}$

$$\left| \frac{1}{\varepsilon} d(\delta_\varepsilon^x v, \delta_\varepsilon^x w) - d^{\delta_\varepsilon^x u} \left(\delta_{\varepsilon^{-1}}^{\delta_\varepsilon^x u} \delta_\varepsilon^x v, \delta_{\varepsilon^{-1}}^{\delta_\varepsilon^x u} \delta_\varepsilon^x w \right) \right| \rightarrow 0 \quad (19)$$

as $\varepsilon \rightarrow 0$. By the axioms A1, A3, the function $(x, u, v) \mapsto d^x(u, v)$ is uniformly continuous on compact sets, being an uniform limit of continuous functions. We prove the fact that the "infinitesimal translations are d^x isometries by passing to the limit in the LHS of ((19)) and by using this uniform continuity.

(II.) If for any x the distance d^x is non degenerate then there exists $C > 0$ such that: for any x and u with $d(x, u) \leq C$ there exists a d^x isometry $\Sigma^x(u, \cdot)$ obtained as the limit:

$$\lim_{\varepsilon \rightarrow 0} \Sigma_\varepsilon^x(u, v) = \Sigma^x(u, v)$$

uniformly with respect to x, u, v in compact set.

Indeed, from the step (I.) we know that $\Delta^x(u, \cdot)$ is a d^x isometry. If d^x is non degenerate then $\Delta^x(u, \cdot)$ is invertible. Let $\Sigma^x(u, \cdot)$ be the inverse.

From proposition 7.2 we know that $\Sigma_\varepsilon^x(u, \cdot)$ is the inverse of $\Delta_\varepsilon^x(u, \cdot)$. Therefore

$$\begin{aligned} d^x(\Sigma_\varepsilon^x(u, w), \Sigma^x(u, w)) &= d^x(\Delta^x(u, \Sigma_\varepsilon^x(u, w)), w) = \\ &= d^x(\Delta^x(u, \Sigma_\varepsilon^x(u, w)), \Delta_\varepsilon^x(u, \Sigma_\varepsilon^x(u, w))). \end{aligned}$$

From the uniformity of convergence in step (I.) and the uniformity assumptions in axioms of dilation structures, the conclusion follows.

(III.) We start by proving that $(U(x), \Sigma^x)$ is a local uniform group. The uniformities are induced by the distance d .

Indeed, according to proposition 7.2, we can pass to the limit with $\varepsilon \rightarrow 0$ and define:

$$inv^x(u) = \lim_{\varepsilon \rightarrow 0} \Delta_\varepsilon^x(u, x) = \Delta^x(u, x).$$

From relation (d) , proposition (7.2) we get (after passing to the limit with $\varepsilon \rightarrow 0$)

$$inv^x(inv^x(u)) = u.$$

We shall see that $inv^x(u)$ is the inverse of u . Relation (c), proposition (7.2) gives:

$$\Delta^x(u, v) = \Sigma^x(inv^x(u), v) \tag{20}$$

therefore relations (a), (b) from proposition 7.2 give

$$\Sigma^x(inv^x(u), \Sigma^x(u, v)) = v, \tag{21}$$

$$\Sigma^x(u, \Sigma^x(u, v)) = v. \tag{22}$$

Relation (e) from proposition 7.2 gives

$$\Sigma^x(u, \Sigma^x(v, w)) = \Sigma^x(\Sigma^x(u, v), w) \tag{23}$$

which shows that Σ^x is an associative operation. From (22), (21) we obtain that for any u, v

$$\Sigma^x(\Sigma^x(inv^x(u), u), v) = v, \tag{24}$$

$$\Sigma^x(\Sigma^x(u, inv^x(u)), v) = v. \tag{25}$$

Remark that for any x, v and $\varepsilon \in (0, 1)$ we have $\Sigma^x(x, v) = v$. Therefore x is a neutral element at left for the operation Σ^x . From the definition of inv^x , relation (20) and the fact that inv^x is equal to its inverse, we get that x is an inverse at right too: for any x, v we have

$$\Sigma^x(v, x) = v.$$

Replace now v by x in relations (24), (25) and prove that indeed $inv^x(u)$ is the inverse of u .

We also have to prove that $(U(x), \Sigma^x)$ admits δ^x as dilations. In this reasoning we need the axiom A2 in strong form.

Namely we have to prove that for any $\mu \in (0, 1)$ we have

$$\delta_\mu^x \Sigma^x(u, v) = \Sigma^x(\delta_\mu^x u, \delta_\mu^x v).$$

For this is sufficient to notice that

$$\delta_\mu^x \Delta_{\varepsilon\mu}^x(u, v) = \Delta_\varepsilon^x(\delta_\mu^x u, \delta_\mu^x v)$$

and pass to the limit as $\varepsilon \rightarrow 0$. \square

Definition 7.4 *The (local) conical group $(U(x), \Sigma^x, \delta^x)$ is the tangent space of (X, d, δ) at x (in the sense of dilation structures). We denote it by $T_x(X, d, \delta) = (U(x), \Sigma^x, \delta^x)$, or by $T_x X$ if (d, δ) are clear from the context.*

Compatibility of topologies. The axiom A3 implies that for any $x \in X$ the function d^x is continuous, therefore open sets with respect to d^x are open with respect to d .

If (X, d) is separable and d^x is non degenerate then the uniformities induced by d and d^x are the same. Indeed, let $\{u_n : n \in \mathbb{N}\}$ be a dense set in $U(x)$, with $x_0 = x$. We can embed $(U(x), (\delta^x, \varepsilon))$ isometrically in the separable Banach space l^∞ , for any $\varepsilon \in (0, 1)$, by the function

$$\phi_\varepsilon(u) = \left(\frac{1}{\varepsilon} d(\delta_\varepsilon^x u, \delta_\varepsilon^x x_n) - \frac{1}{\varepsilon} d(\delta_\varepsilon^x x, \delta_\varepsilon^x x_n) \right)_n.$$

A reformulation of the first part of theorem 7.1 is that on compact sets ϕ_ε uniformly converges to the isometric embedding of $(U(x), d^x)$

$$\phi(u) = (d^x(u, x_n) - d^x(x, x_n))_n.$$

Remark that the uniformity induced by (δ, ε) is the same as the uniformity induced by d , and that it is the same induced from the uniformity on l^∞ by the embedding ϕ_ε . We proved that the uniformities induced by d and d^x are the same.

From previous considerations we deduce the following characterization of tangent spaces associated to a dilation structure.

Corollary 7.5 *Let (X, d, δ) be a strong dilation structure with group $\Gamma = (0, +\infty)$. Then for any $x \in X$ the local group $(U(x), \Sigma^x)$ is locally a simply connected Lie group whose Lie algebra admits a positive graduation (a homogeneous group).*

Proof. Indeed, from previous considerations, $(U(x), \Sigma^x)$ is a locally compact group which admits δ^x as a contractive automorphism group (from theorem 7.3). Instead of Siebert proposition 3.9, we need now a version for local groups. Fortunately, theorem 1.1 [20] states that a locally compact, locally connected, contractible (with Siebert' wording) group is locally isomorphic to a contractive Lie group. \square

Straightforward modifications in the proof of the previous theorem allow us to extend some results to length dilation structures.

Theorem 7.6 *Let (X, d, δ) be a length dilation structure. Then:*

- (a) Σ^x is a local group operation on $U(x)$, with x as neutral element and inv^x as the inverse element function; for any $\varepsilon \in (0, 1]$ the dilation δ_ε^x is an automorphism with respect to the group operation;
- (b) the length functional $l^x = l(x, \cdot)$ is invariant with respect to left translations $\Sigma^x(y, \cdot)$, $y \in U(x)$; moreover, for any $\mu \in (0, 1]$ the equality

$$l(x, \delta_\mu^x c) = \mu l(x, c)$$

Proof. Notice that the axiom A4+ is all that we need in order to transform the proof of theorem 10 [4] into a proof of this theorem. Indeed, for this we need the existence of the limits from A4+ and the algebraic relations from theorem 11 [4] which are true only from A0, A1, A2.

If $(\delta_\varepsilon^x y, c) \in \mathcal{L}_\varepsilon(X, d, \delta)$ then $(x, \Sigma_\varepsilon^x(y, \cdot)c) \in \mathcal{L}_\varepsilon(X, d, \delta)$ and moreover

$$l_\varepsilon(\delta_\varepsilon^x y, c) = l_\varepsilon(x, \Sigma_\varepsilon^x(y, \cdot)c)$$

Indeed, this is true because of the equality:

$$\delta^{\delta_\varepsilon^x y} c = \delta_\varepsilon^x \Sigma_\varepsilon^x(y, \cdot)c$$

By passing to the limit with $\varepsilon \rightarrow 0$ and using A3L and A4+ we get

$$l(x, c) = l(x, \Sigma^x(y, \cdot)c)$$

For any $\varepsilon, \mu > 0$ (and sufficiently small) $(x, c) \in \mathcal{L}_{\varepsilon\mu}(X, d, \delta)$ is equivalent with $(x, \delta_\mu^x c) \in \mathcal{L}_\varepsilon(X, d, \delta)$ and moreover:

$$l_\varepsilon(x, \delta_\mu^x c) = \frac{1}{\varepsilon} l_d(\delta_{\varepsilon\mu}^x c) = \mu l_{\varepsilon\mu}(x, c)$$

We pass to the limit with $\varepsilon \rightarrow 0$ and we get the desired equality. \square

7.3 Differentiability with respect to a pair of dilation structures

For any pair of strong dilation structures or length dilation structures there is an associated notion of differentiability (section 7.2 [4]). First we need the definition of a morphism of conical groups.

Definition 7.7 Let (N, δ) and $(M, \bar{\delta})$ be two conical groups. A function $f : N \rightarrow M$ is a conical group morphism if f is a group morphism and for any $\varepsilon > 0$ and $u \in N$ we have $f(\delta_\varepsilon u) = \bar{\delta}_\varepsilon f(u)$.

The definition of the derivative, or differential, with respect to dilations structures follows. In the case of a pair of Carnot groups this is just the definition of the Pansu derivative introduced in [29].

Definition 7.8 Let (X, d, δ) and $(Y, \bar{d}, \bar{\delta})$ be two strong dilation structures or length dilation structures and $f : U \subset X \rightarrow Y$ be a continuous function defined on an open subset of X . The function f is differentiable in $x \in U$ if there exists a conical group morphism $Df(x) : T_x X \rightarrow T_{f(x)} Y$, defined on a neighbourhood of x with values in a neighbourhood of $f(x)$ such that

$$\lim_{\varepsilon \rightarrow 0} \sup \left\{ \frac{1}{\varepsilon} \bar{d} \left(f(\delta_\varepsilon^x u), \bar{\delta}_\varepsilon^{f(x)} Df(x)(u) \right) : d(x, u) \leq \varepsilon \right\} = 0, \quad (26)$$

The morphism $Df(x)$ is called the derivative, or differential, of f at x .

7.4 Equivalent dilation structures

In the following we adopt a notion of (local) equivalence of dilation structures.

Definition 7.9 *Two strong dilation structures (X, δ, d) and $(X, \bar{\delta}, \bar{d})$ are (locally) equivalent if*

(a) *the identity map $id : (X, d) \rightarrow (X, \bar{d})$ is bilipschitz, uniformly on compact sets, that is for any compact set $K \subset X$ there are numbers $R = R(K) > 0$ and $c = c(K), C = C(K) > 0$ such that for any $x \in K$ the restriction of the identity on the ball $B_d(x, R)$ is bilipschitz, with Lipschitz constant smaller than C and Lipschitz constant of the inverse smaller than by $\frac{1}{c}$,*

(b) *for any $x \in X$ there are functions P^x, Q^x (defined for $u \in X$ sufficiently close to x) such that*

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \bar{d} \left(\delta_\varepsilon^x u, \bar{\delta}_\varepsilon^x Q^x(u) \right) = 0, \quad (27)$$

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} d \left(\bar{\delta}_\varepsilon^x u, \delta_\varepsilon^x P^x(u) \right) = 0, \quad (28)$$

uniformly with respect to x, u in compact sets.

We shall keep the word "local" further, only if needed.

Proposition 7.10 *(X, d, δ) and $(X, \bar{d}, \bar{\delta})$ are equivalent if and only if*

(a) *the identity map $id : (X, d) \rightarrow (X, \bar{d})$ is locally bilipschitz,*

(b) *for any $x \in X$ there are conical group morphisms:*

$$P^x : T_x(X, \bar{\delta}, \bar{d}) \rightarrow T_x(X, \delta, d) \text{ and } Q^x : T_x(X, \delta, d) \rightarrow T_x(X, \bar{\delta}, \bar{d})$$

such that the following limits exist

$$\lim_{\varepsilon \rightarrow 0} \left(\bar{\delta}_\varepsilon^x \right)^{-1} \delta_\varepsilon^x(u) = Q^x(u), \quad (29)$$

$$\lim_{\varepsilon \rightarrow 0} \left(\delta_\varepsilon^x \right)^{-1} \bar{\delta}_\varepsilon^x(u) = P^x(u), \quad (30)$$

and are uniform with respect to x, u in compact sets.

The next theorem shows a link between the tangent bundles of equivalent dilation structures.

Theorem 7.11 *Let (X, d, δ) and $(X, \bar{d}, \bar{\delta})$ be equivalent strong dilation structures. Then for any $x \in X$ and any $u, v \in X$ sufficiently close to x we have:*

$$\bar{\Sigma}^x(u, v) = Q^x \left(\Sigma^x(P^x(u), P^x(v)) \right). \quad (31)$$

The two tangent bundles are therefore isomorphic in a natural sense.

7.5 Distribution of a dilation structure

Let (X, d, δ) be a strong dilation structure or a length dilation structure. We have then a notion of differentiability for curves in X , seen as continuous functions from (a open interval in) \mathbb{R} , with the usual dilation structure, to X with the dilation structure (X, d, δ) . Further we want to see what differentiability in the sense of definition 7.8 means for curves. In proposition 7.13 we shall arrive to a notion of a distribution in a dilation structure, with the geometrical meaning of a cone of all possible derivatives of curves passing through a point.

Definition 7.12 *In a normed conical group N we shall denote by $D(N)$ the set of all $u \in N$ with the property that $\varepsilon \in ((0, \infty), +) \mapsto \delta_\varepsilon u \in N$ is a morphism of groups.*

$D(N)$ is always non empty, because it contains the neutral element of N . $D(N)$ is also a cone, with dilations δ_ε , and a closed set.

Proposition 7.13 *Let (X, d, δ) be a strong dilation structure or a length dilation structure and let $c : [a, b] \rightarrow (X, d)$ be a continuous curve. For any $x \in X$ and any $y \in T_x(X, d, \delta)$ we denote by*

$$\text{inv}^x(y) = \Delta^x(y, x)$$

the inverse of y with respect to the group operation in $T_x(X, d, \delta)$. Then the following are equivalent:

- (a) *c is derivable in $t \in (a, b)$ with respect to the dilation structure (X, d, δ) ;*
- (b) *there exists $\dot{c}(t) \in D(T_{c(t)}(X, d, \delta))$ such that*

$$\begin{aligned} \frac{1}{\varepsilon} d(c(t + \varepsilon), \delta_\varepsilon^{c(t)} \dot{c}(t)) &\rightarrow 0 \\ \frac{1}{\varepsilon} d(c(t - \varepsilon), \delta_\varepsilon^{c(t)} \text{inv}^{c(t)}(\dot{c}(t))) &\rightarrow 0 \end{aligned}$$

Proof. It is straightforward that a conical group morphism $f : \mathbb{R} \rightarrow N$ is defined by its value $f(1) \in N$. Indeed, for any $a > 0$ we have $f(a) = \delta_a f(1)$ and for any $a < 0$ we have $f(a) = \delta_a f(1)^{-1}$. From the morphism property we also deduce that

$$\delta v = \{ \delta_a v : a > 0, v = f(1) \text{ or } v = f(1)^{-1} \}$$

is a one parameter group and that for all $\alpha, \beta > 0$ we have $\delta_{\alpha+\beta} u = \delta_\alpha u \delta_\beta u$. We have therefore a bijection between conical group morphisms $f : \mathbb{R} \rightarrow (N, \delta)$ and elements of $D(N)$.

The curve $c : [a, b] \rightarrow (X, d)$ is derivable in $t \in (a, b)$ if and only if there is a morphism of normed conical groups $f : \mathbb{R} \rightarrow T_{c(t)}(X, d, \delta)$ such that for any $a \in \mathbb{R}$ we have

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} d(c(t + \varepsilon a), \delta_\varepsilon^{c(t)} f(a)) = 0$$

Take $\dot{c}(t) = f(1)$. Then $\dot{c}(t) \in D(T_{c(t)}(X, d, \delta))$. For any $a > 0$ we have $f(a) = \delta_a^{c(t)} \dot{c}(t)$; otherwise if $a < 0$ we have $f(a) = \delta_a^{c(t)} \text{inv}^{c(t)} \dot{c}(t)$. This implies the equivalence stated on the proposition. \square

8 Supplementary properties of dilation structures

At this level of generality, dilation structures come in many flavors. Further we shall propose two supplementary properties which may be satisfied by a dilation structure: the Radon-Nikodym property and the property of being tempered. It will turn out that sub-riemannian spaces may be endowed with dilation structures having the RNP (therefore true in particular for riemannian spaces), but genuinely sub-riemannian spaces don't have dilation structures which are tempered, in contradistinction to the riemannian spaces.

8.1 The Radon-Nikodym property

Definition 8.1 *A strong dilation structure or a length dilation structure (X, d, δ) has the Radon-Nikodym property (or rectifiability property, or RNP) if any Lipschitz curve $c : [a, b] \rightarrow (X, d)$ is derivable almost everywhere.*

Three examples. The first example is obvious. For $(X, d) = (\mathbb{V}, d)$, a real, finite dimensional, normed vector space, with distance d induced by the norm, the (usual) dilations δ_ε^x are given by:

$$\delta_\varepsilon^x y = x + \varepsilon(y - x)$$

Dilations are defined everywhere. Axioms 0,1,2 are obviously true. Concerning the axiom A3, remark that for any $\varepsilon > 0$, $x, u, v \in X$ we have $\frac{1}{\varepsilon}d(\delta_\varepsilon^x u, \delta_\varepsilon^x v) = d(u, v)$. It follows that $d^x = d$ for any $x \in X$. Concerning the axiom A4, remark that for any $\varepsilon > 0$ and any $x, u, v \in X$ we have

$$\Delta_\varepsilon^x(u, v) = x + \varepsilon(u - x) + \frac{1}{\varepsilon}(x + \varepsilon(v - x) - x - \varepsilon(u - x)) = x + \varepsilon(u - x) + v - u$$

therefore this quantity converges to $x + v - u = x + (v - x) - (u - x)$. as $\varepsilon \rightarrow 0$. For this dilation structure, the RNP as in definition 8.1 is just the usual Radon-Nikodym property.

Further is an example of a dilation structure which does not have the Radon-Nikodym property. Take $X = \mathbb{R}^2$ with the euclidean distance d . For any $z = 1 + i\theta \in \mathbb{C}$, with $\theta \in \mathbb{R}$, we define dilations

$$\delta_\varepsilon x = \varepsilon^z x .$$

Then $(\mathbb{R}^2, d, \delta)$ is a dilation structure, with dilations $\delta_\varepsilon^x y = x + \delta_\varepsilon(y - x)$.

Two such dilation structures (constructed with the help of complex numbers $1 + i\theta$ and $1 + i\theta'$) are equivalent if and only if $\theta = \theta'$. Moreover, if $\theta \neq \theta'$ then there are no non trivial Lipschitz curves in X which are differentiable almost everywhere. It means that such a dilation structure does not have the Radon-Nikodym property.

More than this, such a dilation structure does not satisfy the obviously reformulated Rademacher theorem (which states that a Lipschitz function is derivable – in the sense of dilation structures – almost everywhere with respect to the 2-Hausdorff measure). Indeed any holomorphic and Lipschitz function from X to X (holomorphic in the usual sense on $X = \mathbb{R}^2 = \mathbb{C}$) is differentiable almost everywhere (classically and, equivalently, in the sense of dilation structures), but there are Lipschitz functions from X to X which are not differentiable almost everywhere: it suffices to take a C^∞ function from \mathbb{R}^2 to \mathbb{R}^2 which is not holomorphic.

The last example concerns riemannian manifolds endowed with dilation structures as in proposition 5.2.

Proposition 8.2 *The dilation structure of a riemannian manifold constructed from the geodesic spray, as in proposition 5.2, has the RNP from definition 8.1.*

Proof. Indeed, locally this dilation structure is trivially equivalent to the first dilation structure which was constructed in section 5.3. That dilation structure is just the one of \mathbb{R}^n with the usual dilations and an euclidean distance. The RNP in this case is just the usual RNP, therefore, by corollary 8.5 (from the section 8.2), we get the result. \square

8.2 Radon-Nikodym property, representation of length, distributions

Theorem 8.3 *Let (X, d, δ) be a strong dilation structure with the Radon-Nikodym property, over a complete length metric space (X, d) . Then for any $x, y \in X$ we have*

$$d(x, y) = \inf \left\{ \int_a^b d^{c(t)}(c(t), \dot{c}(t)) dt : c : [a, b] \rightarrow X \text{ Lipschitz}, \right. \\ \left. c(a) = x, c(b) = y \right\}$$

Proof. By theorem 2.15, for almost every $t \in (a, b)$ the upper dilation of c in t can be expressed as the limit

$$Lip(c)(t) = \lim_{s \rightarrow t} \frac{d(c(s), c(t))}{|s - t|}$$

For a dilation structure with the RNP, for almost every $t \in [a, b]$ there is $\dot{c}(t) \in D(T_{c(t)}X)$ such that

$$\frac{1}{\varepsilon} d(c(t + \varepsilon), \delta_\varepsilon^{c(t)} \dot{c}(t)) \rightarrow 0$$

It follows that for almost every $t \in [a, b]$ we have

$$Lip(c)(t) = \lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} d(c(t + \varepsilon), c(t)) = d^{c(t)}(c(t), \dot{c}(t))$$

which implies the representation of length. \square

As a consequence, the distance d is uniquely determined by the distribution in the sense of dilation structures and the norm on it.

Corollary 8.4 *Let (X, d, δ) and $(X, \bar{d}, \bar{\delta})$ be two strong dilation structures with the Radon-Nikodym property, which are also complete length metric spaces, such that for any $x \in X$ we have $D(T_x(X, d, \delta)) = D(T_x(X, \bar{d}, \bar{\delta}))$ and $d^x(x, u) = \bar{d}^x(x, u)$ for any $u \in D(T_x(X, d, \delta))$. Then $d = \bar{d}$.*

Another consequence is that the RNP is transported by the equivalence of dilation structures. More precisely we have the following.

Corollary 8.5 *Let (X, d, δ) and $(X, \bar{d}, \bar{\delta})$ be equivalent strong dilation structures. Then for any $x \in X$ we have*

$$Q^x(D(T_x(X, \delta, d))) = D(T_x(X, \bar{\delta}, \bar{d}))$$

If (X, d, δ) has the Radon-Nikodym property, then $(X, \bar{d}, \bar{\delta})$ has the same property.

Suppose that (X, d, δ) and $(X, \bar{d}, \bar{\delta})$ are complete length spaces with the Radon-Nikodym property. If the functions P^x, Q^x from definition 7.9 (b) are isometries, then $d = \bar{d}$.

8.3 Tempered dilation structures

The notion of a tempered dilation structure extends the results of Venturini [31] and Buttazzo, De Pascale and Fragala [16] (propositions 2.3, 2.6 and a part of theorem 3.1) to dilation structures.

The following definition associates a class of distances $\mathcal{D}(X, \bar{d}, \bar{\delta})$ to a strong dilation structure $(X, \bar{d}, \bar{\delta})$. This is a which generalization of the class of distances $\mathcal{D}(X)$ from [16], definition 2.1.

Definition 8.6 *To a strong dilation structure $(X, \bar{d}, \bar{\delta})$ we associate the class $\mathcal{D}(X, \bar{d}, \bar{\delta})$ of all length distance functions d on X such that for any $\varepsilon > 0$ and any x, u, v sufficiently close there are constants $0 < c < C$ with the property*

$$c \bar{d}^x(u, v) \leq \frac{1}{\varepsilon} d(\bar{\delta}_\varepsilon^x u, \bar{\delta}_\varepsilon^x v) \leq C \bar{d}^x(u, v) \quad (32)$$

The dilation structure $(X, \bar{d}, \bar{\delta})$ is tempered if $\bar{d} \in \mathcal{D}(X, \bar{d}, \bar{\delta})$.

On $\mathcal{D}(X, \bar{d}, \bar{\delta})$ we put the topology of uniform convergence (induced by distance \bar{d}) on compact subsets of $X \times X$.

To any distance $d \in \mathcal{D}(X, \bar{d}, \bar{\delta})$ we associate the function:

$$\phi_d(x, u) = \limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} d(x, \delta_\varepsilon^x u)$$

defined for any $x, u \in X$ sufficiently close. We have therefore

$$c \bar{d}^x(x, u) \leq \phi_d(x, u) \leq C \bar{d}^x(x, u) \quad (33)$$

Notice that if $d \in \mathcal{D}(X, \bar{d}, \bar{\delta})$ then for any x, u, v sufficiently close we have the relations

$$\begin{aligned} & -\bar{d}(x, u) O(\bar{d}(x, u)) + c \bar{d}^x(u, v) \leq \\ & \leq d(u, v) \leq C \bar{d}^x(u, v) + \bar{d}(x, u) O(\bar{d}(x, u)) \end{aligned}$$

Important remark. If $c : [0, 1] \rightarrow X$ is a d -Lipschitz curve and $d \in \mathcal{D}(X, \bar{d}, \bar{\delta})$ then we may decompose it in a finite family of curves c_1, \dots, c_n (with n depending on c) such that there are $x_1, \dots, x_n \in X$ with c_k is \bar{d}^{x_k} -Lipschitz. Indeed, the image of the curve $c([0, 1])$ is compact, therefore we may cover it with a finite number of balls $B(c(t_k), \rho_k, \bar{d}^{c(t_k)})$ and apply (32). If moreover $(X, \bar{d}, \bar{\delta})$ is tempered then it follows that $c : [0, 1] \rightarrow X$ d -Lipschitz curve is equivalent with c \bar{d} -Lipschitz curve.

By using the same arguments as in the proof of theorem 8.3, we get the following extension of proposition 2.4 [16].

Proposition 8.7 *If $(X, \bar{d}, \bar{\delta})$ is tempered, with the Radon-Nikodym property, and $d \in \mathcal{D}(X, \bar{d}, \bar{\delta})$ then*

$$\begin{aligned} d(x, y) = \inf \left\{ \int_a^b \phi_d(c(t), \dot{c}(t)) dt : c : [a, b] \rightarrow X \text{ } \bar{d}\text{-Lipschitz,} \right. \\ \left. c(a) = x, c(b) = y \right\} \end{aligned}$$

The next theorem is a generalization of a part of theorem 3.1 [16].

Theorem 8.8 *Let $(X, \bar{d}, \bar{\delta})$ be a strong dilation structure which is tempered, with the Radon-Nikodym property, and $d_n \in \mathcal{D}(X, \bar{d}, \bar{\delta})$ a sequence of distances converging to $d \in \mathcal{D}(X, \bar{d}, \bar{\delta})$. Denote by L_n, L the length functional induced by the distance d_n , respectively by d . Then L_n Γ -converges to L .*

Proof. We have to prove the liminf inequality and the existence of a recovery sequence, i.e. parts (a), (b) respectively, of the definition 6.2 of Γ -convergence of length functionals. The proof is inspired by the one of implication (i) \Rightarrow (iii) from theorem 3.1 [16], p. 252-253, we only need to replace everywhere expressions like $|x - y|$ by $\bar{d}(x, y)$ and use proposition 8.7, relations (33) and (32) instead of respectively proposition 2.4 and relations (2.6) and (2.3) [16].

Proof of (a). Let us take any sequence of curves $(c_n)_n$, with d_n -Lipschitz curve $c_n : [0, 1] \rightarrow X$ for every n . We suppose that c_n converges uniformly to the curve c . We want to prove that

$$L(c) \leq \liminf_{n \rightarrow \infty} L_n(c_n)$$

For any $\eta > 0$ there is a number $m = m(\eta)$ and a division $\Delta_\eta = \{t_0 = 0, \dots, t_m = 1\}$ of $[0, 1]$ such that

$$L(c) - \eta \leq \sum_{i=0}^{m-1} d(c(t_i), c(t_{i+1}))$$

From the uniform convergence of c_n to c it follows that there is a compact set $K \subset X \times X$ and a number $N = N(K)$ such that for any $n \geq N$ and any $s, t \in [0, 1]$ we have $(c_n(s), c_n(t)) \in K$.

For any $n \geq N(K)$,

$$\begin{aligned} |d_n(c_n(t_i), c_n(t_{i+1})) - d(c(t_i), c(t_{i+1}))| &\leq |d_n(c_n(t_i), c_n(t_{i+1})) - d(c_n(t_i), c_n(t_{i+1}))| + \\ &+ |d(c_n(t_i), c_n(t_{i+1})) - d(c(t_i), c(t_{i+1}))| \leq \sup_K |d_n - d| + 2 \sup_{[0,1]} d(c_n, c) \leq \varepsilon_n \end{aligned}$$

where $\varepsilon_n = \sup_K |d_n - d| + 2C \sup_{[0,1]} \bar{d}d(c_n, c)$ and the number $C > 0$ comes from $d \in \mathcal{D}(X, \bar{d}, \bar{\delta})$, see the comments after definition 8.6. From this inequality we obtain:

$$L(c) \leq \eta + L_n(c_n) + \varepsilon_n m(\eta)$$

We pass to the limit with $n \rightarrow \infty$ and we use $\varepsilon_n \rightarrow 0$ (from the convergence of c_n to c) and we get:

$$L(c) \leq \eta + \liminf_{n \rightarrow \infty} L_n(c_n)$$

which gives the desired liminf inequality by the fact that η is arbitrary.

Proof of (b). We have the curve c which is d -Lipschitz (therefore \bar{d} -Lipschitz) and we want to construct a recovery sequence of curves c_n .

Let us consider an increasing sequence $k(n)$ of natural numbers, for the moment without any supplementary assumption. For each n we division $[0, 1]$ into $k(n)$ intervals of equal length, denote by t_i^n the elements of the division, $i = 0, \dots, k(n)$, and we define the curve c_n to be one such that $c_n(t_i^n) = c(t_i^n)$ for all i and, denoting by c_n^i the restriction of c_n to the interval $[t_i^n, t_{i+1}^n]$, such that c_n^i to be an almost geodesic with respect to d_n , that is

$$L_n(c_n^i) \leq d_n(c(t_i^n), c(t_{i+1}^n)) + \frac{1}{2^{k(n)}} \quad (34)$$

We want to prove that c_n is a recovery sequence, for a well chosen sequence $k(n)$.

It is not restricting the generality to suppose that the length $L(c)$ is equal to one. Let $K \subset X \times X$ be a compact set which contains the set $\{(c(t), y) : d(c(t), y) \leq 1\}$. We choose then the sequence $k(n)$ to be one with the property

$$\lim_{n \rightarrow \infty} k(n) \sup_K |d_n - d| = 0$$

Then, there is a number $N(K)$ such that for any $n \geq N(K)$ and any For any $t \in [0, 1]$ and any $s, t \in [0, 1]$ we have $(c_n(s), c_n(t)) \in K$.

For any $n \geq N(K)$ and any $t \in [0, 1]$ we denote by $[t_n^-, t_n^+]$ the interval of the division of $[0, 1]$ into $k(n)$ intervals of equal length, where t belongs. Let also c_n^t be the restriction of c_n to the interval $[t_n^-, t_n^+]$.

Then $\sup_{[0,1]} \bar{d}(c_n(t), c(t)) \leq A_n + B_n$, where $A_n = \sup_{[0,1]} \bar{d}(c(t_n^+), c(t))$ and $B_n = \sup_{[0,1]} \bar{d}(c_n(t_n^+), c_n(t))$. Trivially $A_n \rightarrow 0$ as $n \rightarrow \infty$. The term B_n may be estimated as follows. By proposition 8.7, relations (33), (32) and (34), there are numbers $0 < c < C$ such that

$$c \bar{d}(c_n(t_n^+), c_n(t)) \leq L_n(c_n^t) + \frac{1}{2^{k(n)}} \leq C \bar{d}(c(t_n^-), c(t_n^+)) + \frac{1}{2^{k(n)}}$$

We have then

$$\begin{aligned} L(c) &\leq \sum_{i=0}^{k(n)-1} d(c(t_i^n), c(t_{i+1}^n)) = \sum_{i=0}^{k(n)-1} d_n(c_n(t_i^n), c_n(t_{i+1}^n)) + \\ &+ \sum_{i=0}^{k(n)-1} (d(c(t_i^n), c(t_{i+1}^n)) - d_n(c(t_i^n), c(t_{i+1}^n))) \geq L_n(c_n) - \frac{k(n)}{2^{k(n)}} - k(n) \sup_K |d - d_n| \end{aligned}$$

We apply $\limsup_{n \rightarrow \infty}$ to his inequality and we prove the claim. \square

As a corollary we obtain a large class of examples of length dilation structures.

Corollary 8.9 *If $(X, \bar{d}, \bar{\delta})$ is a strong dilation structure which is tempered and it has the Radon-Nikodym property then it is a length dilation structure.*

Proof. Indeed, from the hypothesis we deduce that $\bar{\delta}_\varepsilon \bar{d} \in \mathcal{D}(X, \bar{d}, \bar{\delta})$. For any sequence $\varepsilon_n \rightarrow 0$ we thus obtain a sequence of distances $d_n = \bar{\delta}_{\varepsilon_n} \bar{d}$ converging to \bar{d}^x . We apply now theorem 8.8 and we get the result. \square

We arrive therefore to the following characterization of riemannian manifolds.

Theorem 8.10 *The dilation structure associated to a riemannian manifold, as in proposition 5.2, is tempered and is a length dilation structure.*

Conversely, if (X, d, δ) is a strong dilation structure which is tempered, it has the Radon-Nikodym property and moreover for any $x \in X$ the tangent space is a commutative local group, then any open, with compact closure subset of X can be endowed with a C^1 riemannian structure which gives a distance d' which is bilipschitz equivalent with d .

Proof. We already know that this dilation structure has the RNP. It is also tempered, because of the estimate (9) from the proof of the proposition 2.22. By corollary 8.9 it is a length dilation structure.

For the converse assertion, remark that the only local conical groups, which are locally compact and admit a one parameter group of dilations (that is the abelian group Γ is $(0, +\infty)$ are (open neighbourhoods of 0 in) \mathbb{R}^n . From the fact that the dilation structure is tempered, it follows that locally (X, d) is bilipschitz with \mathbb{R}^n endowed with the norm constructed as (33). But any norm on \mathbb{R}^n is bilipschitz with an euclidean norm. \square

9 Dilation structures on sub-riemannian manifolds

In [6], followed in this section, we proved that we can associate dilation structures to regular sub-Riemannian manifolds. This result is the source of inspiration of the notion of a coherent projection, section 10.1.

In this section we use differential geometric tools, mainly the normal frames, definition 9.7. This has been proved by Bellaïche [3], starting with theorem 4.15 and ending in the first half of section 7.3 (page 62). We shall not give an exposition of this long proof, although a streamlined version of it would be very useful.

From the existence of normal frames we shall get the existence of certain dilation structures regular sub-riemannian manifolds, theorem 9.9. From this, according to the general theory of dilation structures, via Siebert type results also, follow all classical results concerning the structure of the tangent space to a point of a regular sub-riemannian manifold.

In particular, this is evidence for the fact that the classical differential calculus is needed only in the part concerning the existence of normal frames and after this stage an intrinsic way of reasoning is possible.

Let us compare with, maybe, the closest results, which are those of Vodopyanov [32], [33]. In both approaches the tangent space to a point is defined only locally, as a neighbourhood of the point, in the manifold. The difference is that Vodopyanov proves the existence of the (locally defined) operation on the tangent space under very weak regularity assumptions on the sub-riemannian manifold, by using the differential structure of the underlying manifold. In distinction, we prove in [4], in an abstract setting, that the very existence of a dilation structure induces a locally defined operation.

9.1 Sub-riemannian manifolds

M is a connected, n dimensional, real manifold. A (differential geometric) distribution on M is a smooth subbundle D of M . Such a distribution associates to any point $x \in M$ a vector space $D_x \subset T_x M$. The dimension of the distribution D at point $x \in M$ is $m(x) = \dim D_x$. The function $x \in M \mapsto m(x)$ is locally constant, because of the distribution is smooth. We shall suppose that the dimension of the distribution is globally constant and we denote it by m . In general $m \leq n$. The typical case we are interested in is $m < n$.

A horizontal curve $c : [a, b] \rightarrow M$ is a curve which is almost everywhere derivable and for almost any $t \in [a, b]$ we have $\dot{c}(t) \in D_{c(t)}$. The class of horizontal curves will be denoted by $Hor(M, D)$.

Definition 9.1 *The distribution D is completely non-integrable if M is locally connected by horizontal curves, that is curves in $Hor(M, D)$.*

The Chow condition (C) [17] is sufficient for the distribution D to be completely non-integrable.

Theorem 9.2 (Chow) *Let us suppose there is a positive integer number k (called the rank of the distribution D) such that for any $x \in X$ there is a topological open ball $U(x) \subset M$ with $x \in U(x)$ such that there are smooth vector fields X_1, \dots, X_m in $U(x)$ with the property:*

(C) *the vector fields X_1, \dots, X_m span D_x and these vector fields together with their iterated brackets of order at most k span the tangent space $T_y M$ at every point $y \in U(x)$.*

Then the distribution D is completely non-integrable in the sense of definition 9.1.

Definition 9.3 *A sub-riemannian (SR) manifold is a triple (M, D, g) , where M is a connected manifold, D is a completely non-integrable distribution on M , and g is a metric (Euclidean inner-product) on the distribution (or horizontal bundle) D .*

With the notations from condition (C), let us define on $U = U(x)$ a filtration of bundles as follows. First we define the class of horizontal vector fields on U

$$\mathcal{X}^1(U(x), D) = \{X \in \mathcal{X}^\infty(U) : \forall y \in U(x), X(y) \in D_y\}$$

Next, we define inductively for all positive integers j the following vector fields:

$$\mathcal{X}^{j+1}(U(x), D) = \mathcal{X}^j(U(x), D) + [\mathcal{X}^1(U(x), D), \mathcal{X}^j(U(x), D)]$$

where $[\cdot, \cdot]$ denotes the bracket of vector fields. All in all we obtain the filtration $\mathcal{X}^j(U(x), D) \subset \mathcal{X}^{j+1}(U(x), D)$. By evaluation at $y \in U(x)$, we get a filtration

$$V^j(y, U(x), D) = \{X(y) : X \in \mathcal{X}^j(U(x), D)\}$$

According to Chow condition (C), there is a positive integer k such that for all $y \in U(x)$ we have

$$D_y = V^1(y, U(x), D) \subset V^2(y, U(x), D) \subset \dots \subset V^k(y, U(x), D) = T_y M$$

Consequently, to the sub-riemannian manifold is associated the string of numbers:

$$\nu_1(y) = \dim V^1(y, U(x), D) < \nu_2(y) = \dim V^2(y, U(x), D) < \dots < n = \dim M$$

Generally $k, \nu_j(y)$ may vary from a point to another. The number k is called the step of the distribution at y .

Definition 9.4 *The distribution D , which satisfies the Chow condition (C), is regular if $\nu_j(y)$ are constant on the manifold M .*

The sub-riemannian manifold (M, D, g) is regular if D is regular and for any $x \in M$ there is a topological ball $U(x) \subset M$ with $x \in U(x)$ and an orthonormal (with respect to the metric g) family of smooth vector fields $\{X_1, \dots, X_m\}$ in $U(x)$ which satisfy the condition (C).

The length of a horizontal curve is obtained from the metric g by

$$l(c) = \int_a^b (g_{c(t)}(\dot{c}(t), \dot{c}(t)))^{\frac{1}{2}} dt$$

Definition 9.5 *The Carnot-Carathéodory distance (or CC distance) associated to the sub-riemannian manifold is the distance induced by the length l of horizontal curves:*

$$d(x, y) = \inf \{l(c) : c \in \text{Hor}(M, D), c(a) = x, c(b) = y\}$$

The Chow condition ensures the existence of a horizontal path linking any two sufficiently closed points, therefore the CC distance is locally finite. The distance depends only on the distribution D and metric g , and not on the choice of vector fields X_1, \dots, X_m satisfying the condition (C). The space (M, d) is locally compact and complete, and the topology induced by the distance d is the same as the topology of the manifold M . (These important details may be recovered from reading carefully the proofs of Chow theorem given by Bellaïche [3] or Gromov [25].)

9.2 Sub-riemannian dilation structures associated to normal frames

In the following we suppose that M is a regular sub-riemannian manifold. The dimension of M as a differential manifold is denoted by n , the step of the regular sub-riemannian manifold (M, D, g) is denoted by k , the dimension of the distribution is m , and there are numbers $\nu_j, j = 1, \dots, k$ such that for any $x \in M$ we have $\dim V^j(x) = \nu_j$. The Carnot-Carathéodory distance is denoted by d . Further, we stay in a small open neighbourhood of an arbitrary, but fixed point $x_0 \in M$. We shall no longer mention the dependence of various objects on x_0 , on the neighbourhood $U(x_0)$, or the distribution D .

Definition 9.6 An adapted frame $\{X_1, \dots, X_n\}$ is an ordered collection of smooth vector fields, constructed according to the following recipe.

The first m vector fields X_1, \dots, X_m satisfy the condition (C). Therefore, for any point x the vectors $X_1(x), \dots, X_m(x)$ form a basis for D_x .

We associate to any word $a_1 \dots a_q$ with letters in the alphabet $1, \dots, m$ the vector field equal to the multi-bracket $[X_{a_1}, [\dots, X_{a_q}]]$. We add, in the lexicographic order, $n - m$ elements to the set $\{X_1, \dots, X_m\}$ until we get a collection $\{X_1, \dots, X_n\}$ such that: for any $j = 1, \dots, k$ and for any point x the set $\{X_1(x), \dots, X_{\nu_j}(x)\}$ is a basis for $V^j(x)$.

Let $\{X_1, \dots, X_n\}$ be an adapted frame. For any $j = 1, \dots, n$ the degree $\deg X_j$ of the vector field X_j is defined as the only positive integer p such that for any point x we have

$$X_j(x) \in V_x^p \setminus V_x^{p-1}(x)$$

In definition below, the key are the uniform convergence assumptions. This is in line with Gromov suggestions in the last section of Bellaïche [3].

Definition 9.7 A normal frame is an adapted frame $\{X_1, \dots, X_n\}$ which satisfies the following supplementary properties:

(a) the limit

$$\lim_{\varepsilon \rightarrow 0^+} \frac{1}{\varepsilon} d \left(\exp \left(\sum_1^n \varepsilon^{\deg X_i} a_i X_i \right) (y), y \right) = A(y, a) \in (0, +\infty)$$

exists and is uniform with respect to y in compact sets and $a = (a_1, \dots, a_n) \in W$, with $W \subset \mathbb{R}^n$ compact neighbourhood of $0 \in \mathbb{R}^n$,

(b) for any compact set $K \subset M$ with diameter (with respect to the CC distance d) sufficiently small, and for any $i = 1, \dots, n$ there are functions

$$P_i(\cdot, \cdot, \cdot) : U_K \times U_K \times K \rightarrow \mathbb{R}$$

with $U_K \subset \mathbb{R}^n$ a sufficiently small compact neighbourhood of $0 \in \mathbb{R}^n$ such that for any $x \in K$ and any $a, b \in U_K$ we have

$$\exp \left(\sum_1^n a_i X_i \right) (x) = \exp \left(\sum_1^n P_i(a, b, y) X_i \right) \circ \exp \left(\sum_1^n b_i X_i \right) (x)$$

and such that the following limit exists

$$\lim_{\varepsilon \rightarrow 0^+} \varepsilon^{-\deg X_i} P_i(\varepsilon^{\deg X_j} a_j, \varepsilon^{\deg X_k} b_k, x) \in \mathbb{R}$$

and it is uniform with respect to $x \in K$ and $a, b \in U_K$.

The condition (a) definition 9.7 is a part of the conclusion of Gromov approximation theorem, namely when one point coincides with the center of nilpotentization; also condition (b) is equivalent with a statement of Gromov concerning the convergence of rescaled vector fields to their nilpotentization (an informed reader must at least follow in all details the papers Bellaïche [3] and Gromov [25], where differential calculus in the classical sense is heavily used).

In the case of a Lie group G endowed with a left invariant distribution, normal frames are very easy to visualize. The left-invariant distribution is completely non-integrable if and only it is generated by the left translation of a vector subspace D of the algebra $\mathfrak{g} = T_e G$ which generates the whole Lie algebra \mathfrak{g} . Let us take $\{X_1, \dots, X_m\}$ to be a collection of $m = \dim D$ left-invariant independent vector fields. Let us define with their help an adapted frame, as explained in definition 9.6. This frame is in fact normal.

Definition 9.8 Let (M, d, g) be a regular sub-riemannian manifold and let $\{X_1, \dots, X_n\}$ be a normal frame (locally defined, but for simplicity here we neglect this detail). To this pair we associate a triple (M, d, δ) , where: d is the Carnot-Carathéodory distance, and for any point $x \in M$ and any $\varepsilon \in (0, +\infty)$ (sufficiently small if necessary), the dilation δ_ε^x is defined by the expression

$$\delta_\varepsilon^x \left(\exp \left(\sum_{i=1}^n a_i X_i \right) (x) \right) = \exp \left(\sum_{i=1}^n a_i \varepsilon^{\deg X_i} X_i \right) (x)$$

We shall prove that (M, d, δ) is a dilation structure. This allows us to get the main results concerning the infinitesimal geometry of a regular sub-riemannian manifold.

Theorem 9.9 Let (M, D, g) be a regular sub-riemannian manifold and $U \subset M$ an open set which admits a normal frame. Then (U, d, δ) is a strong dilation structure.

Proof. We only need to prove that A3 and A4 hold. The fact that A3 is true is a result similar to Gromov local approximation theorem [25], p. 135, or to Bellaïche theorem 7.32 [3].

A3 is satisfied. We prove that the limit $\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} d(\delta_\varepsilon^x u, \delta_\varepsilon^x v) = d^x(u, v)$ exists and it uniform with respect to x, u, v sufficiently close. We represent u and v with respect to the normal frame:

$$u = \exp \left(\sum_1^n u_i X_i \right) (x) \quad , \quad v = \exp \left(\sum_1^n v_i X_i \right) (x)$$

Let us denote by $u_\varepsilon = \delta_\varepsilon^x u = \exp \left(\sum_1^n \varepsilon^{\deg X_i} u_i X_i \right) (x)$. Then

$$\begin{aligned} \frac{1}{\varepsilon} d(\delta_\varepsilon^x u, \delta_\varepsilon^x v) &= \frac{1}{\varepsilon} d \left(\delta_\varepsilon^x \exp \left(\sum_1^n u_i X_i \right) (x), \delta_\varepsilon^x \exp \left(\sum_1^n v_i X_i \right) (x) \right) = \\ &= \frac{1}{\varepsilon} d \left(\exp \left(\sum_1^n \varepsilon^{\deg X_i} u_i X_i \right) (x), \exp \left(\sum_1^n \varepsilon^{\deg X_i} v_i X_i \right) (x) \right) \end{aligned}$$

Let us make the notation: for any $i = 1, \dots, n$ $a_i^\varepsilon = \varepsilon^{-\deg X_i} P_i(\varepsilon^{\deg X_j} v_j, \varepsilon^{\deg X_k} u_k, x)$. By the second part of property (b), definition 9.7, the vector $a^\varepsilon \in \mathbb{R}^n$ converges to a finite value $a^0 \in \mathbb{R}^n$, as $\varepsilon \rightarrow 0$, uniformly with respect to x, u, v in compact set. In the same time u_ε converges to x , as $\varepsilon \rightarrow 0$. But remark that

$$\begin{aligned} \frac{1}{\varepsilon} d(\delta_\varepsilon^x u, \delta_\varepsilon^x v) &= \frac{1}{\varepsilon} d \left(u_\varepsilon, \exp \left(\sum_1^n P_i(\varepsilon^{\deg X_j} v_j, \varepsilon^{\deg X_k} u_k, x) X_i \right) (u_\varepsilon) \right) = \\ &= \frac{1}{\varepsilon} d \left(u_\varepsilon, \exp \left(\sum_1^n \varepsilon^{\deg X_i} (\varepsilon^{-\deg X_i} P_i(\varepsilon^{\deg X_j} v_j, \varepsilon^{\deg X_k} u_k, x)) X_i \right) (u_\varepsilon) \right) = \\ &= \frac{1}{\varepsilon} d \left(u_\varepsilon, \exp \left(\sum_1^n \varepsilon^{\deg X_i} a_i^\varepsilon X_i \right) (u_\varepsilon) \right) \end{aligned}$$

Now we use the uniform convergence assumptions from definition 9.7. For fixed $\eta > 0$ the term

$$B(\eta, \varepsilon) = \frac{1}{\varepsilon} d \left(u_\eta, \exp \left(\sum_1^n \varepsilon^{\deg X_i} a_i^\eta X_i \right) (u_\eta) \right)$$

converges to a real number $A(u_\eta, a_\eta)$ as $\varepsilon \rightarrow 0$, uniformly with respect to u_η and a_η . Since u_η converges to x and a_η converges to a^0 as $\eta \rightarrow 0$, by the uniform convergence assumption in (a), definition 9.7 we get that

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} d(\delta_\varepsilon^x u, \delta_\varepsilon^x v) = \lim_{\eta \rightarrow 0} A(u_\eta, a_\eta) = A(x, a^0)$$

A4 is satisfied. As ε converges to 0, the approximate difference $\Delta_\varepsilon^x(u, v)$ has a limit, which is uniform with respect to x, u, v sufficiently close. Indeed,

$$\Delta_\varepsilon^x(u, v) = \delta_{\varepsilon^{-1}}^{u_\varepsilon} \exp \left(\sum_1^n \varepsilon^{\deg X_i} v_i X_i \right) (x)$$

We use the first part of the property (b), definition 9.7, in order to write

$$\exp \left(\sum_1^n \varepsilon^{\deg X_i} v_i X_i \right) (x) = \exp \left(\sum_1^n P_i(\varepsilon^{\deg X_j} v_j, \varepsilon^{\deg X_k} u_k, x) X_i \right) (u_\varepsilon)$$

We finish the computation:

$$\Delta_\varepsilon^x(u, v) = \exp \left(\sum_1^n \varepsilon^{-\deg X_i} P_i(\varepsilon^{\deg X_j} v_j, \varepsilon^{\deg X_k} u_k, x) X_i \right) (u_\varepsilon)$$

As ε goes to 0 the point u_ε converges to x uniformly with respect to x, u sufficiently close (as a corollary of the previous theorem, for example). The proof therefore ends by invoking the second part of the property (b), definition 9.7. \square

10 Coherent projections: a dilation structure looks down on another

The equivalence of dilation structures, definition 7.9, may be seen as a composite of two partial order relations.

Definition 10.1 *A strong dilation structure (X, δ, d) is looking down on another strong dilation structure $(X, \bar{\delta}, \bar{d})$ if*

- (a) *the identity map $id : (X, d) \rightarrow (X, \bar{d})$ is lipschitz and*
- (b) *for any $x \in X$ there are functions Q^x (defined for $u \in X$ sufficiently close to x) such that*

$$\lim_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon} \bar{d} \left(\delta_\varepsilon^x u, \bar{\delta}_\varepsilon^x Q^x(u) \right) = 0, \quad (35)$$

uniformly with respect to x, u in compact sets.

This leads us to the introduction of coherent projections, i.e. to the study of pairs of dilation structures, one looking down on another. (The name is inspired by the notion of a set looking down on another introduced in [19].) We prefer to work with a pair $(\bar{\delta}, Q)$ made by a dilation structure and a function Q as in the previous definition, point (b), instead of working with a pair of dilation structures.

10.1 Coherent projections

Definition 10.2 Let $(X, \bar{d}, \bar{\delta})$ be a strong dilation structure. A coherent projection of $(X, \bar{d}, \bar{\delta})$ is a function which associates to any $x \in X$ and $\varepsilon \in (0, 1]$ a map $Q_\varepsilon^x : U(x) \rightarrow X$ such that:

(I) $Q_\varepsilon^x : U(x) \rightarrow Q_\varepsilon^x(U(x))$ is invertible and the inverse is $Q_{\varepsilon^{-1}}^x$; moreover, the following commutation relation holds: for any $\varepsilon, \mu > 0$ and any $x \in X$

$$Q_\varepsilon^x \bar{\delta}_\mu^x = \bar{\delta}_\mu^x Q_\varepsilon^x$$

(II) the limit $\lim_{\varepsilon \rightarrow 0} Q_\varepsilon^x u = Q^x u$ is uniform with respect to x, u in compact sets.

(III) for any $\varepsilon, \mu > 0$ and any $x \in X$ we have $Q_\varepsilon^x Q_\mu^x = Q_{\varepsilon\mu}^x$. Also $Q_1^x = id$ and $Q_\varepsilon^x x = x$.

(IV) define $\Theta_\varepsilon^x(u, v) = \bar{\delta}_{\varepsilon^{-1}}^x Q_{\varepsilon^{-1}}^{\bar{\delta}_\varepsilon^x Q_\varepsilon^x u} \bar{\delta}_\varepsilon^x Q_\varepsilon^x v$. Then the limit exists

$$\lim_{\varepsilon \rightarrow 0} \Theta_\varepsilon^x(u, v) = \Theta^x(u, v)$$

and it is uniform with respect to x, u, v in compact sets.

A coherent projection induces the dilations $\delta_\varepsilon^x = \bar{\delta}_\varepsilon^x Q_\varepsilon^x$.

Proposition 10.3 Let $(X, \bar{d}, \bar{\delta})$ be a strong dilation structure and Q a coherent projection. Then:

(a) the induced dilations commute with the old ones: for any $\varepsilon, \mu > 0$ and any $x \in X$ we have $\delta_\varepsilon^x \bar{\delta}_\mu^x = \bar{\delta}_\mu^x \delta_\varepsilon^x$.

(b) as $\varepsilon \rightarrow 0$ the coherent projection becomes a projection: for any $x \in X$ we have $Q^x Q^x = Q^x$.

(c) the induced dilations δ satisfy the conditions A1, A2, A4 from definition 4.4,

(d) the following relations are true (we denote by $\Sigma^x, \Delta^x, \dots$, the approximate or infinitesimal sum and difference computed with the help of induced dilations):

$$\Theta_\varepsilon^x(u, v) = \bar{\Sigma}_\varepsilon^x(Q_\varepsilon^x u, \Delta_\varepsilon^x(u, v)) \quad (36)$$

$$\Delta^x(u, v) = \bar{\Delta}^x(Q^x u, \Theta^x(u, v)) \quad (37)$$

$$Q^x \Delta^x(u, v) = \bar{\Delta}^x(Q^x u, Q^x v) \quad (38)$$

Proof. (a) this is a consequence of the commutativity condition (I) (second part). Indeed, we have $\delta_\varepsilon^x \bar{\delta}_\mu^x = \bar{\delta}_\varepsilon^x Q_\varepsilon^x \bar{\delta}_\mu^x = \bar{\delta}_\varepsilon^x \bar{\delta}_\mu^x Q_\varepsilon^x = \bar{\delta}_\mu^x \bar{\delta}_\varepsilon^x Q_\varepsilon^x = \bar{\delta}_\mu^x \delta_\varepsilon^x$.

(b) we pass to the limit $\varepsilon \rightarrow 0$ in the equality $Q_{\varepsilon^2}^x = Q_\varepsilon^x Q_\varepsilon^x$ and we get, based on condition (II), that $Q^x Q^x = Q^x$.

(c) Axiom A1 for δ is equivalent with (III). Indeed, the equality $\delta_\varepsilon^x \bar{\delta}_\mu^x = \bar{\delta}_{\varepsilon\mu}^x$ is equivalent with: $\bar{\delta}_{\varepsilon\mu}^x Q_{\varepsilon\mu}^x = \bar{\delta}_{\varepsilon\mu}^x Q_\varepsilon^x Q_\mu^x$. This is true because $Q_\varepsilon^x Q_\mu^x = Q_{\varepsilon\mu}^x$. We also have $\delta_1^x = \bar{\delta}_1^x Q_1^x = Q_1^x = id$. Moreover $\delta_\varepsilon^x x = \bar{\delta}_\varepsilon^x Q_\varepsilon^x x = Q_\varepsilon^x \bar{\delta}_\varepsilon^x x = Q_\varepsilon^x x = x$. Let us compute now:

$$\Delta_\varepsilon^x(u, v) = \delta_{\varepsilon^{-1}}^{\delta_\varepsilon^x u} \delta_\varepsilon^x v = \bar{\delta}_{\varepsilon^{-1}}^{\delta_\varepsilon^x u} Q_{\varepsilon^{-1}}^{\delta_\varepsilon^x u} \delta_\varepsilon^x v = \bar{\delta}_{\varepsilon^{-1}}^{\delta_\varepsilon^x u} \bar{\delta}_\varepsilon^x \Theta_\varepsilon^x(u, v) = \bar{\Delta}_\varepsilon^x(Q_\varepsilon^x u, \Theta_\varepsilon^x(u, v))$$

We can pass to the limit in the last term of this string of equalities and we prove that the axiom A4 is satisfied by δ : there exists the limit

$$\Delta^x(u, v) = \lim_{\varepsilon \rightarrow 0} \Delta_\varepsilon^x(u, v) \quad (39)$$

which is uniform as written in A4, moreover we have the equality (36).

(d) We pass to the limit to the limit with $\varepsilon \rightarrow 0$ in the relation (36) and we obtain (37). In order to prove (38) we notice that:

$$Q_{\varepsilon}^{\delta_{\varepsilon}^x u} \Delta_{\varepsilon}^x(u, v) = Q_{\varepsilon}^{\delta_{\varepsilon}^x u} \delta_{\varepsilon-1}^{\delta_{\varepsilon}^x u} \delta_{\varepsilon}^x v = \bar{\delta}_{\varepsilon-1}^{\delta_{\varepsilon}^x u} \bar{\delta}_{\varepsilon}^x Q_{\varepsilon}^x v = \bar{\Delta}_{\varepsilon}^x(Q_{\varepsilon}^x u, Q_{\varepsilon}^x v)$$

which gives(38) by passing to the limit with $\varepsilon \rightarrow 0$ in this relation. \square

Induced dilations at a scale. For any $x \in X$ and $\varepsilon \in (0, 1)$ the dilation δ_{ε}^x can be seen as an isomorphism of strong dilation structures with coherent projections:

$$\delta_{\varepsilon}^x : (U(x), \delta_{\varepsilon}^x \bar{d}, \hat{\delta}_{\varepsilon}^x, \hat{Q}_{\varepsilon}^x) \rightarrow (\delta_{\varepsilon}^x U(x), \frac{1}{\varepsilon} \bar{d}, \bar{\delta}, Q)$$

We may use this morphism in order to transport the induced dilations and the coherent projection:

$$\hat{\delta}_{\varepsilon, \mu}^{x, u} = \delta_{\varepsilon-1}^x \bar{\delta}_{\mu}^{\delta_{\varepsilon}^x u} \delta_{\varepsilon}^x \quad , \quad \hat{Q}_{\varepsilon, \mu}^{x, u} = \delta_{\varepsilon-1}^x Q_{\mu}^{\delta_{\varepsilon}^x u} \delta_{\varepsilon}^x$$

The dilation $\bar{\delta}_{\varepsilon}^x$, which is an isomorphism of strong dilation structures with coherent projections:

$$\bar{\delta}_{\varepsilon}^x : (U(x), \bar{\delta}_{\varepsilon}^x \bar{d}, \bar{\delta}_{\varepsilon}^x, \bar{Q}_{\varepsilon}^x) \rightarrow (\bar{\delta}_{\varepsilon}^x U(x), \frac{1}{\varepsilon} \bar{d}, \bar{\delta}, Q)$$

may be also used to transport induced dilations and coherent projections:

$$\bar{\delta}_{\varepsilon, \mu}^{x, u} = \bar{\delta}_{\varepsilon-1}^x \bar{\delta}_{\mu}^{\bar{\delta}_{\varepsilon}^x u} \bar{\delta}_{\varepsilon}^x \quad , \quad \bar{Q}_{\varepsilon, \mu}^{x, u} = \bar{\delta}_{\varepsilon-1}^x \bar{Q}_{\mu}^{\bar{\delta}_{\varepsilon}^x u} \bar{\delta}_{\varepsilon}^x$$

These two morphisms are related: because $\delta_{\varepsilon}^x = \bar{\delta}_{\varepsilon}^x Q_{\varepsilon}^x$ it follows that

$$Q_{\varepsilon}^x : (U(x), \delta_{\varepsilon}^x \bar{d}, \hat{\delta}_{\varepsilon}^x, \hat{Q}_{\varepsilon}^x) \rightarrow (Q_{\varepsilon}^x U(x), \bar{\delta}_{\varepsilon}^x \bar{d}, \bar{\delta}_{\varepsilon}^x, \bar{Q}_{\varepsilon}^x)$$

is yet another isomorphism of strong dilation structures with coherent projections. This isomorphism measures the "distortion" between the previous two isomorphisms. Recall that in the limit Q_{ε}^x becomes a projection, therefore $Q_{\varepsilon}^x U(x)$ will be squeezed to a flattened version $Q^x U(x)$ (which will represent, in the case of sub-riemannian manifolds, a local version of the distribution evaluated at x).

We shall denote the derivative of a curve with respect to the dilations $\hat{\delta}_{\varepsilon}^x$ by $\frac{\hat{d}_{\varepsilon}^x}{dt}$. Also, the derivative of the curve c with respect to $\bar{\delta}$ is denoted by $\frac{\bar{d}}{dt}$. A curve c is $\hat{\delta}_{\varepsilon}^x$ -derivable if and only if $\delta_{\varepsilon}^x c$ is $\bar{\delta}$ -derivable and

$$\frac{\hat{d}_{\varepsilon}^x}{dt} c(t) = \delta_{\varepsilon-1}^x \frac{\bar{d}}{dt} (\delta_{\varepsilon}^x c)(t)$$

Q-horizontal curves. These will play an important role further, here is the definition.

Definition 10.4 *Let $(X, \bar{d}, \bar{\delta})$ be a strong dilation structure and Q a coherent projection. A curve $c : [a, b] \rightarrow X$ is Q -horizontal if for almost any $t \in [a, b]$ the curve c is derivable and the derivative of c at t , denoted by $\dot{c}(t)$ has the property:*

$$Q^{c(t)} \dot{c}(t) = \dot{c}(t) \tag{40}$$

A curve $c : [a, b] \rightarrow X$ is Q -everywhere horizontal if for all $t \in [a, b]$ the curve c is derivable and the derivative has the horizontality property (40).

If the curve $\delta_\varepsilon^x c$ is Q -horizontal then $\frac{\bar{d}_\varepsilon^x}{dt}(Q_\varepsilon^x c)(t) = \Theta_\varepsilon^x(c(t), \frac{\hat{d}_\varepsilon^x}{dt}c(t))$. Indeed,

$$\frac{\bar{d}_\varepsilon^x}{dt}(Q_\varepsilon^x c)(t) = \bar{\delta}_{\varepsilon^{-1}}^x Q^{\delta_\varepsilon^x c(t)} \bar{\delta}_\varepsilon^x \frac{\bar{d}_\varepsilon^x}{dt}(Q_\varepsilon^x c)(t)$$

which implies: $\bar{\delta}_\varepsilon^x \frac{\bar{d}_\varepsilon^x}{dt}(Q_\varepsilon^x c)(t) = Q^{\delta_\varepsilon^x c(t)} \bar{\delta}_{\varepsilon^{-1}}^x \frac{\bar{d}_\varepsilon^x}{dt}(Q_\varepsilon^x c)(t) = Q^{\delta_\varepsilon^x c(t)} \bar{\delta}_\varepsilon^x \frac{\hat{d}_\varepsilon^x}{dt}c(t)$.

10.2 Length functionals associated to coherent projections

Definition 10.5 Let $(X, \bar{d}, \bar{\delta})$ be a strong dilation structure with the Radon-Nikodym property and Q a coherent projection. We define the associated distance $d : X \times X \rightarrow [0, +\infty]$ by:

$$d(x, y) = \inf \left\{ \int_a^b \bar{d}^{c(t)}(c(t), \dot{c}(t)) dt : c : [a, b] \rightarrow X \text{ } \bar{d}\text{-Lipschitz, } \right. \\ \left. c(a) = x, c(b) = y, \text{ and } \forall a.e. t \in [a, b] \quad Q^{c(t)} \dot{c}(t) = \dot{c}(t) \right\}$$

As usual, we accept that the distance between two points may be infinite. The equivalence relation $x \equiv y$ if $d(x, y) < +\infty$ induces a decomposition of the space X into a reunion of equivalence classes, each equivalence class being a set connected by horizontal curves with finite length. Later on we shall give a sufficient condition (the generalized Chow condition (Cgen)) on the coherent projection Q for X to be (locally) connected by horizontal curves.

If X is connected by horizontal curves and (X, d) is complete then d is a length distance. For proving this, it is sufficient to check that d has the approximate middle property: for any $\varepsilon > 0$ and for any $x, y \in X$ there exists $z \in X$ such that

$$\max \{d(x, z), d(y, z)\} \leq \frac{1}{2} d(x, y) + \varepsilon$$

For any $\varepsilon > 0$ there exists a horizontal curve $c : [a, b] \rightarrow X$ such that $c(a) = x, c(b) = y$ and $d(x, y) + 2\varepsilon \geq l(c)$ (where $l(c)$ is the length of c with respect to the distance \bar{d}). There is then a $\tau \in [a, b]$ such that

$$\int_a^\tau \bar{d}^{c(t)}(c(t), \dot{c}(t)) dt = \int_\tau^b \bar{d}^{c(t)}(c(t), \dot{c}(t)) dt = \frac{1}{2} l(c)$$

Let now $z = c(\tau)$. We have then: $\max \{d(x, z), d(y, z)\} \leq \frac{1}{2} l(c) \leq \frac{1}{2} d(x, y) + \varepsilon$, which proves the claim.

Notations concerning length functionals. The length functional associated to the distance \bar{d} is denoted by \bar{l} . In the same way the length functional associated with $\bar{\delta}_\varepsilon^x$ is denoted by \bar{l}_ε^x .

We introduce the space $\mathcal{L}_\varepsilon(X, d, \delta) \subset X \times Lip([0, 1], X, d)$:

$$\mathcal{L}_\varepsilon(X, d, \delta) = \{(x, c) \in X \times \mathcal{C}([0, 1], X) : c : [0, 1] \in U(x), \\ \delta_\varepsilon^x c \text{ is } \bar{d}\text{-Lip, } Q\text{-horizontal and } Lip(\delta_\varepsilon^x c) \leq 2\varepsilon l_d(\delta_\varepsilon^x c)\}$$

For any $\varepsilon \in (0, 1)$ we define the length functional

$$l_\varepsilon : \mathcal{L}_\varepsilon(X, d, \delta) \rightarrow [0, +\infty] \quad , \quad l_\varepsilon(x, c) = l_\varepsilon^x(c) = \frac{1}{\varepsilon} \bar{l}(\delta_\varepsilon^x c)$$

By theorem 8.3 we have:

$$l_\varepsilon^x(c) = \int_0^1 \frac{1}{\varepsilon} \bar{d}^{\delta_\varepsilon^x c(t)} \left(\delta_\varepsilon^x c(t), \frac{\bar{d}}{dt}(\delta_\varepsilon^x c)(t) \right) dt = \int_0^1 \frac{1}{\varepsilon} \bar{d}^{\delta_\varepsilon^x c(t)} \left(\delta_\varepsilon^x c(t), \delta_\varepsilon^x \frac{\hat{d}_\varepsilon^x}{dt}c(t) \right) dt$$

10.3 Conditions (A) and (B)

Further are two supplementary hypotheses on a coherent projection Q .

Definition 10.6 *Let $(X, \bar{d}, \bar{\delta})$ be a strong dilation structure, Q a coherent projection and δ the induced dilation.*

(A) δ_ε^x is \bar{d} -bilipschitz in compact sets in the following sense: for any compact set $K \subset X$ and for any $\varepsilon \in (0, 1]$ there is a number $L(K) > 0$ such that for any $x \in K$ and u, v sufficiently close to x we have:

$$\frac{1}{\varepsilon} \bar{d}(\delta_\varepsilon^x u, \delta_\varepsilon^x v) \leq L(K) \bar{d}(u, v)$$

(B) if $u = Q^x u$ then the curve $t \in [0, 1] \mapsto Q^x \delta_t^x u = \bar{\delta}_t^x u = \delta_t^x u$ is Q -everywhere horizontal and for any $a \in [0, 1]$ we have

$$\limsup_{a \rightarrow 0} \frac{\bar{l}(t \in [0, a] \mapsto \bar{\delta}_t^x u)}{\bar{d}(x, \bar{\delta}_a^x u)} = 1$$

uniformly with respect to x, u in compact set K .

The condition (A) implies that for any \bar{d} -Lipschitz curve c , the "rescaled" curve $\delta_\varepsilon^x c$ is also Lipschitz. Moreover, the condition (A) is equivalent with the fact that Q_ε^x is locally $\bar{\delta}_\varepsilon^x \bar{d}$ -Lipschitz, where we use the notation

$$(\bar{\delta}_\varepsilon^x \bar{d})(u, v) = \frac{1}{\varepsilon} \bar{d}(\bar{\delta}_\varepsilon^x u, \bar{\delta}_\varepsilon^x v)$$

More precisely, condition (A) is equivalent with: for any compact set $K \subset X$ and for any $\varepsilon \in (0, 1]$ there is a number $L(K) > 0$ such that for any $x \in K$ and u, v sufficiently close to x we have:

$$(\bar{\delta}_\varepsilon^x \bar{d})(Q_\varepsilon^x u, Q_\varepsilon^x v) \leq L(K) \bar{d}(u, v) \quad (41)$$

Indeed, we have:

$$(\bar{\delta}_\varepsilon^x \bar{d})(Q_\varepsilon^x u, Q_\varepsilon^x v) = \frac{1}{\varepsilon} \bar{d}(\delta_\varepsilon^x u, \delta_\varepsilon^x v) \leq L(K) \bar{d}(u, v)$$

The condition (B) is a generalization of the fact that the "distribution" $x \mapsto Q^x U(x)$ is generated by horizontal one parameter flows, see theorem 11.1.

11 Distributions in sub-riemannian spaces as coherent projections

Here we explain how coherent projections appear in sub-riemannian geometry.

Let $\{Y_1, \dots, Y_n\}$ be a frame induced by a parameterization $\phi : O \subset \mathbb{R}^n \rightarrow U \subset M$ of a small open, connected set U in the manifold M . This parameterization induces a dilation structure on U , by

$$\tilde{\delta}_\varepsilon^{\phi(a)} \phi(b) = \phi(a + \varepsilon(-a + b))$$

We take the distance $\tilde{d}(\phi(a), \phi(b)) = \|b - a\|$.

Let $\{X_1, \dots, X_n\}$ be a normal frame, cf. definition 9.7, let d be the Carnot-Carathéodory distance and let

$$\delta_\varepsilon^x \left(\exp \left(\sum_{i=1}^n a_i X_i \right) (x) \right) = \exp \left(\sum_{i=1}^n a_i \varepsilon^{deg X_i} X_i \right) (x)$$

be the dilation structure associated, by theorem 9.9.

Alternatively, we may take another dilation structure, constructed as follows: extend the metric g on the distribution D to a riemannian metric on M , denoted for convenience also by g . Let \bar{d} be the riemannian distance induced by the riemannian metric g , and the dilations

$$\bar{\delta}_\varepsilon^x \left(\exp \left(\sum_{i=1}^n a_i X_i \right) (x) \right) = \exp \left(\sum_{i=1}^n a_i \varepsilon X_i \right) (x)$$

Then $(U, \bar{d}, \bar{\delta})$ is a strong dilation structure which is equivalent with the dilation structure $(U, \bar{d}, \bar{\delta})$.

From now on we may define coherent projections associated either to the pair $(\tilde{\delta}, \delta)$ or to the pair $(\bar{\delta}, \delta)$.

Let us define Q_ε^x by:

$$Q_\varepsilon^x \left(\exp \left(\sum_{i=1}^n a_i X_i \right) (x) \right) = \exp \left(\sum_{i=1}^n a_i \varepsilon^{\deg X_i - 1} X_i \right) (x) \quad (42)$$

Theorem 11.1 *Q is a coherent projection associated with the dilation structure $(U, \bar{d}, \bar{\delta})$ which satisfies conditions (A) and (B) definition 10.6. Moreover, for any point x the image of Q^x is in the "distribution" $D(T_x(U, d, \delta))$ (from proposition 7.13), where (U, d, δ) is the strong dilation structure from theorem 9.9.*

Proof. (I) definition 10.2 is true, because $\delta_\varepsilon^x u = Q_\varepsilon^x \bar{\delta}_\varepsilon^x$ and $\delta_\varepsilon^x \bar{\delta}_\varepsilon^x = \bar{\delta}_\varepsilon^x \delta_\varepsilon^x$. (II), (III) and (IV) are consequences of these facts and theorem 9.9, with a proof similar to the one of proposition 10.3.

Definition (42) of the coherent projection Q implies that:

$$Q^x \left(\exp \left(\sum_{i=1}^n a_i X_i \right) (x) \right) = \exp \left(\sum_{\deg X_i = 1} a_i X_i \right) (x) \quad (43)$$

The projection Q^x has the property: for any x and

$$u = \exp \left(\sum_{\deg X_i = 1} a_i X_i \right) (x)$$

we have $Q^x u = u$ and the curve

$$s \in [0, 1] \mapsto \delta_s^x u = \exp \left(s \sum_{\deg X_i = 1} a_i X_i \right) (x)$$

is D -horizontal and joins x and u . This implies condition (B). As for the condition (A), according to the comments after definition 10.6, it just means that the coherent projection given by (42) is locally Lipschitz (with a Lipschitz constant uniform with respect to x in compact set) with respect to the rescaled riemannian distance $\bar{\delta}_\varepsilon^x \bar{d}$. But the dilation structure $(U, \bar{d}, \bar{\delta})$ is tempered, according to theorem 8.10, therefore the rescaled distance is bilipshitz with the riemannian distance \bar{d} , again with Lipschitz constants which are uniform with respect to x in compact set. From (42) we easily get that Q_ε^x is (uniformly w.r.t. x) Lipschitz w.r.t. the riemannian distance \bar{d} . In conclusion (A) is true.

After examination of the construction of the dilation structure (U, d, δ) from theorem 9.9, we see that $D(T_x(U, d, \delta))$ (from proposition 7.13) is exactly the set of all points (sufficiently close to x)

which can be written as exponentials of vectors in the (classical differential) geometric distribution. Therefore the last statement of the theorem is proved. \square

We may equally define a coherent projection which induces the dilations δ from $\tilde{\delta}$. Also, if we change the chosen normal frame with another of the same kind, then we pass to a dilation structure which is equivalent to (U, d, δ) . In conclusion, coherent projections are not geometrical objects per se, but in a natural way one may define a notion of equivalent coherent projections such that the equivalence class is geometrical, i.e. independent of the choice of a pair of particular dilation structures, each in a given equivalence class. Another way of putting this is that a class of equivalent dilation structures may be seen as a category and a coherent projection is a functor between such categories. We shall not pursue this line here. Anyway, the only advantage of choosing $\tilde{\delta}, \delta$ related by the normal frame $\{X_1, \dots, X_n\}$ is that they are associated with a coherent projection with a simple expression.

12 An intrinsic description of sub-riemannian geometry

12.1 The generalized Chow condition

We want to transform the Chow condition (C), theorem 9.2, into a statement formulated in terms of coherent projections. Essentially, the Chow condition (C) tells us that the (sub-riemannian) space is locally connected by horizontal curves which are constructed from concatenations of exponentials of horizontal vector fields.

In the following we need to explain what are the correspondents of exponentials of horizontal vector fields, then we need a way to manage the concatenation procedure. A simple explanation is this:

- (a) the exponentials of horizontal vector fields are, approximately, the induced dilations of the coherent projection,
- (b) A concatenation of those exponentials is coded by a word made by letters, each letters coding one exponential.

A supplementary complication is that we need to have an efficient way to manage these concatenations of exponentials *at any scale*. That is why we start with a scale, that is with a $\varepsilon > 0$ and with the coherent projection, dilations, and so on, at that scale (i.e. those transported by some properly chosen dilation field).

We shall follow closely [7] section 10, getting into details as necessary.

Words over an alphabet. This is a standard way of notation. We shall need words as codes for curves, as explained previously. For any "alphabet" (that is a set) A we denote by A^* the collection of finite words $q = a_1 \dots a_p$, $p \in \mathbb{N}$, $p > 0$. The empty word (with no letters) is denoted by \emptyset . The length of the word $q = a_1 \dots a_p$ is $|q| = p$; the length of the empty word is 0. We may need words which are infinite at right. The set of those words over the alphabet A is denoted by A^ω . For any word $w \in A^\omega \cup A^*$ and any $p \in \mathbb{N}$ we denote by $[w]_p$ the finite word obtained from the first p letters of w (if $p = 0$ then $[w]_0 = \emptyset$ (in the case of a finite word q , if $p > |q|$ then $[q]_p = q$).

For any non-empty $q_1, q_2 \in A^*$ and $w \in A^\omega$ the concatenation of q_1 and q_2 is the finite word $q_1 q_2 \in A^*$ and the concatenation of q_1 and w is the (infinite) word $q_1 w \in A^\omega$. For any $q \in A^*$ and $w \in A^\omega$ we have $q \emptyset = q$ (as concatenation of finite words) and $\emptyset w = w$ (as concatenation of a finite empty word and an infinite word).

Words as codes for Q -horizontal curves. To the coherent projection Q of a strong dilation structure $(X, \bar{d}, \bar{\delta})$ we associate a family of transformations, which correspond to concatenations of exponentials of horizontal vector fields, at a given scale.

Definition 12.1 To any word $w \in (0, 1]^\omega$ and any $\varepsilon \in (0, 1]$ we associate the transformation

$$\Psi_{\varepsilon w} : X_{\varepsilon w}^* \subset X^* \setminus \{\emptyset\} \rightarrow X^*$$

defined recursively by the following procedure.

If $w = \emptyset$ then we define $\Psi_{\varepsilon \emptyset}^1(x) = x$. For any non-empty finite word $q = xx_1 \dots x_p \in X_{\varepsilon w}^*$ and for any $k \geq 1$ we have

$$\Psi_{\varepsilon \emptyset}^{k+1}([q]_{k+1}) = \delta_{\varepsilon^{-1}}^x Q^{\delta_\varepsilon^x \Psi_{\varepsilon w}^k([q]_k)} \delta_\varepsilon^x q_{k+1}$$

If w is not the empty word then we define the functions $\Psi_{\varepsilon w}^k$ by: $\Psi_{\varepsilon w}^1(x) = x$, and for any $k \geq 1$ we have

$$\Psi_{\varepsilon w}^{k+1}([q]_{k+1}) = \delta_{\varepsilon^{-1}}^x Q_{w_k}^{\delta_\varepsilon^x \Psi_{\varepsilon w}^k([q]_k)} \delta_\varepsilon^x q_{k+1} \quad (44)$$

Finally, for any non-empty finite word $q = xx_1 \dots x_p \in X_{\varepsilon w}^*$ we put

$$\Psi_{\varepsilon w}(xx_1 \dots x_p) = \Psi_{\varepsilon w}^1(x) \dots \Psi_{\varepsilon w}^{k+1}(xx_1 \dots x_k) \dots \Psi_{\varepsilon w}^{p+1}(xx_1 \dots x_p)$$

The domain $X_{\varepsilon w}^* \subset X^* \setminus \{\emptyset\}$ is such that the previous definitions make sense.

By using the definition of a coherent projection, we can give the following, more precise description of $X_{\varepsilon w}^*$ as follows: for any compact set $K \subset X$ there is $\rho = \rho(K) > 0$ such that for any $x \in K$ the word $q = xx_1 \dots x_p \in X_{\varepsilon w}^*$ if for any $k \geq 1$ we have

$$\bar{d}(x_{k+1}, \Psi_{\varepsilon w}^k([q]_k)) \leq \rho$$

Let us see what this gives for $\varepsilon = 1$. By definition 12.1 for $\varepsilon = 1$ we have:

$$\Psi_{1w}^1(x) = x \quad , \quad \Psi_{1w}^2(x, x_1) = Q_{w_1}^x x_1 \quad , \quad \Psi_{1w}^3(x, x_1, x_2) = Q_{w_2}^{Q_{w_1}^x x_1} x_2 \quad \dots$$

Proposition 12.2 Suppose that the coherent projection Q satisfies the condition (B) and let $y \in X$ be

$$y = \Psi_{1\emptyset}^{k+1}(xx_1 \dots x_k)$$

Then there is a Q -horizontal curve joining x and y .

Proof. By condition (B) the curve $t \in [0, 1] \mapsto \bar{\delta}_t^x Q^x u$ is a Q -horizontal curve which joins x with $Q^x u$.

Therefore by applying repeatedly the condition (B) we get that there is a Q -horizontal curve between $\Psi_{1\emptyset}^k(xx_1 \dots x_{k-1})$ and $\Psi_{1\emptyset}^{k+1}(xx_1 \dots x_k)$ for any $k > 1$ and a Q -horizontal curve joining x and $\Psi_{1\emptyset}^2(xx_1)$. Therefore by concatenation we get the desired curve. \square

The following are algebraic properties of the transformations $\Psi_{\varepsilon w}$, which explain how they behave with respect to scale.

Proposition 12.3 With the notations from definition 12.1 we have:

(a) $\Psi_{\varepsilon w} \Psi_{\varepsilon \emptyset} = \Psi_{\varepsilon \emptyset}$. Therefore we have the equality of sets:

$$\Psi_{\varepsilon \emptyset} (X_{\varepsilon \emptyset}^* \cap xX^*) = \Psi_{\varepsilon w} (\Psi_{\varepsilon \emptyset} (X_{\varepsilon \emptyset}^* \cap xX^*))$$

(b) $\Psi_{\varepsilon \emptyset}^{k+1}(xq_1 \dots q_k) = \delta_{\varepsilon^{-1}}^x \Psi_{1\emptyset}^{k+1}(x\delta_\varepsilon^x q_1 \dots \delta_\varepsilon^x q_k)$

(c) $\lim_{\varepsilon \rightarrow 0} \delta_{\varepsilon^{-1}}^x \Psi_{1\emptyset}^{k+1}(x\delta_\varepsilon^x q_1 \dots \delta_\varepsilon^x q_k) = \Psi_{0\emptyset}^{k+1}(xq_1 \dots q_k)$ uniformly with respect to x, q_1, \dots, q_k in compact set.

Proof. (a) We use induction on k to prove that for any natural number k we have:

$$\Psi_{\varepsilon w}^{k+1}(\Psi_{\varepsilon\emptyset}^1(x)\dots\Psi_{\varepsilon\emptyset}^{k+1}(xq_1\dots q_k)) = \Psi_{\varepsilon\emptyset}^{k+1}(xq_1\dots q_k) \quad (45)$$

For $k = 0$ we have to prove that $x = x$ which is trivial. For $k = 1$ we have to prove that

$$\Psi_{\varepsilon w}^2(\Psi_{\varepsilon\emptyset}^1(x)\Psi_{\varepsilon\emptyset}^2(xq_1)) = \Psi_{\varepsilon\emptyset}^2(xq_1)$$

This means:

$$\Psi_{\varepsilon w}^2(x\delta_{\varepsilon^{-1}}^x Q^x \delta_{\varepsilon}^x q_1) = \delta_{\varepsilon^{-1}}^x Q_{w_1}^x \delta_{\varepsilon}^x \delta_{\varepsilon^{-1}}^x Q^x \delta_{\varepsilon}^x x_1 = \delta_{\varepsilon^{-1}}^x Q^x \delta_{\varepsilon}^x x_1 = \Psi_{\varepsilon\emptyset}^2(xq_1)$$

Suppose now that $l \geq 2$ and for any $k \leq l$ the relations (45) are true. Then, as previously, it is easy to check (45) for $k = l + 1$.

(b) is true by direct computation. The point (c) is a straightforward consequence of (b) and definition of coherent projections. \square

The following definition is quantitative: it says that a point x is "nested" in a neighbourhood U with respect to the parameters N (natural number) and ε (scale) if there is a small ball of radius ρ around x , with respect to the scaled distance $(\bar{\delta}_{\varepsilon}^x \bar{d})$, such that: the ball is inside U and we can connect x with any y in the ball, by using at most N concatenations of curves coming from the transformations introduced in definition 12.1, such that all these curves stay in U .

Definition 12.4 Let $N \in \mathbb{N}$ be a strictly positive natural number and $\varepsilon \in (0, 1]$. We say that $x \in X$ is (ε, N, Q) -nested in a open neighbourhood $U \subset X$ if there is $\rho > 0$ such that for any finite word $q = x_1\dots x_N \in X^N$, if $(\bar{\delta}_{\varepsilon}^x \bar{d})(x_{k+1}, \Psi_{\varepsilon\emptyset}^k([xq]_k)) \leq \rho$ for any $k = 1, \dots, N$ then we have $q \in U^N$.

If $x \in U$ is (ε, N, Q) -nested then denote by $U(x, \varepsilon, N, Q, \rho) \subset U^N$ the collection of words $q \in U^N$ such that $\bar{\delta}_{\varepsilon}^x \bar{d}(x_{k+1}, \Psi_{\varepsilon\emptyset}^k([xq]_k)) < \rho$ for any $k = 1, \dots, N$.

In the next definition we introduce the condition (Cgen). Its effect is that if the coherent projection Q satisfies (A) and (B) then in the space $(U(x), \bar{\delta}_{\varepsilon}^x)$, with coherent projection \hat{Q}_{ε}^x , we can join any two sufficiently close points by a sequence of at most N horizontal curves. Moreover there is a control on the length of these curves via condition (B) and condition (Cgen).

The function F which appears in the next definition is not specified in general. In the case of sub-riemannian geometry the function F can be taken as $F(\eta) = C\eta^{1/m}$ with m positive natural number (the step of the distribution). Compare with the Folland-Stein lemma 14.

Definition 12.5 A coherent projection Q satisfies the generalized Chow condition if:

(Cgen) for any compact set K there are $\rho = \rho(K) > 0$, $r = r(K) > 0$, a natural number $N = N(Q, K)$ and a function $F(\eta) = \mathcal{O}(\eta)$ such that for any $x \in K$ and $\varepsilon \in (0, 1]$ there are neighbourhoods $U(x), V(x)$ such that any $x \in K$ is (ε, N, Q) -nested in $U(x)$, $B(x, r, \bar{\delta}_{\varepsilon}^x \bar{d}) \subset V(x)$ and such that the mapping

$$x_1\dots x_N \in U(x, N, Q, \rho) \mapsto \Psi_{\varepsilon\emptyset}^{N+1}(x_1\dots x_N)$$

is surjective from $U(x, \varepsilon, N, Q, \rho)$ to $V(x)$. Moreover for any $z \in V(x)$ there exist $y_1, \dots, y_N \in U(x, \varepsilon, N, Q, \rho)$ such that $z = \Psi_{\varepsilon\emptyset}^{N+1}(xy_1, \dots, y_N)$ and for any $k = 0, \dots, N - 1$ we have

$$\bar{\delta}_{\varepsilon}^x \bar{d}(\Psi_{\varepsilon\emptyset}^{k+1}(xy_1\dots y_k), \Psi_{\varepsilon\emptyset}^{k+2}(xy_1\dots y_{k+1})) \leq F(\bar{\delta}_{\varepsilon}^x \bar{d}(x, z))$$

Suppose now that the coherent projection Q satisfies conditions (A), (B) and (Cgen) and let us consider $\varepsilon \in (0, 1]$ and $x, y \in K$, K compact in X . Then there are numbers $N = N(Q, K)$,

$\rho = \rho(Q, K) > 0$ and words $x_1 \dots x_N \in U(x, \varepsilon, N, Q, \rho)$ such that $y = \Psi_{\varepsilon 0}^{N+1}(xx_1 \dots x_N)$. To these data we associate a "short curve" $c : [0, N] \rightarrow X$, which joins x and y , by: for any $t \in [0, N]$

$$c(t) = \bar{\delta}_{\varepsilon, t+N-k}^{x, \Psi_{\varepsilon 0}^{k+1}(xx_1 \dots x_k)} Q_{\Psi_{\varepsilon 0}^{k+1}(xx_1 \dots x_k)} x_{k+1}$$

By extension, any increasing linear reparameterization of a curve c like the one described previously, will be called "short curve" as well.

12.2 The candidate tangent space

Let $(X, \bar{d}, \bar{\delta})$ be a strong dilation structure and Q a coherent projection. Then we have the induced dilations and coherent projections

$$\bar{\delta}_{\mu}^{x, u} v = \Sigma^x(u, \delta_{\mu}^x \Delta^x(u, v)) \quad , \quad \bar{Q}_{\mu}^{x, u} v = \Sigma^x(u, Q_{\mu}^x \Delta^x(u, v))$$

For any curve $c : [0, 1] \rightarrow U(x)$ which is $\bar{\delta}^x$ -derivable and \bar{Q}^x -horizontal almost everywhere:

$$\frac{\bar{d}^x}{dt} c(t) = \bar{Q}^{x, u} \frac{d^x}{dt} c(t)$$

we define the length

$$l^x(c) = \int_0^1 \bar{d}^x \left(x, \Delta^x(c(t), \frac{d^x}{dt} c(t)) \right) dt$$

and the distance function:

$$\bar{d}^x(u, v) = \inf \left\{ l^x(c) : c : [0, 1] \rightarrow U(x) \text{ is } \bar{\delta}^x\text{-derivable,} \right. \\ \left. \text{and } \bar{Q}^x\text{-horizontal a.e., } c(0) = u, c(1) = v \right\}$$

We want to prove that $(U(x), \bar{d}^x, \bar{\delta}^x)$ is a strong dilation structure and \bar{Q}^x is a coherent projection. For this we need first the following theorem.

Theorem 12.6 *The curve $c : [0, 1] \rightarrow U(x)$ is $\bar{\delta}^x$ -derivable, \bar{Q}^x -horizontal almost everywhere, and $l^x(c) < +\infty$ if and only if the curve $Q^x c$ is $\bar{\delta}^x$ -derivable almost everywhere and $\bar{l}^x(Q^x c) < +\infty$. Moreover, we have $\bar{l}^x(Q^x c) = l^x(c)$.*

Proof. The curve c is \bar{Q}^x -horizontal almost everywhere if and only if for almost any $t \in [0, 1]$ we have

$$Q^x \Delta^x(c(t), \frac{d^x}{dt} c(t)) = \Delta^x(c(t), \frac{d^x}{dt} c(t))$$

We shall prove that c is \bar{Q}^x -horizontal is equivalent with

$$\Theta^x(c(t), \frac{d^x}{dt} c(t)) = \frac{\bar{d}^x}{dt} (Q^x c)(t) \quad (46)$$

Indeed, (46) is equivalent with $\lim_{\varepsilon \rightarrow 0} \bar{\delta}_{\varepsilon-1}^x \bar{\Delta}^x(Q^x c(t), Q^x c(t+\varepsilon)) = \bar{\Delta}^x(Q^x c(t), \Theta^x(c(t), \frac{d^x}{dt} c(t)))$, which

is equivalent with $\lim_{\varepsilon \rightarrow 0} \bar{\delta}_{\varepsilon-1}^x \bar{\Delta}^x(Q^x c(t), Q^x c(t+\varepsilon)) = \Delta^x(c(t), \frac{d^x}{dt} c(t))$, finally equivalent with:

$$\lim_{\varepsilon \rightarrow 0} \bar{\delta}_{\varepsilon-1}^x \bar{\Delta}^x(Q^x c(t), Q^x c(t+\varepsilon)) = \lim_{\varepsilon \rightarrow 0} \delta_{\varepsilon-1}^x \Delta^x(c(t), c(t+\varepsilon)) \quad (47)$$

The horizontality condition for the curve c can be written as:

$$\lim_{\varepsilon \rightarrow 0} Q^x \delta_{\varepsilon^{-1}}^x \Delta^x(c(t), c(t + \varepsilon)) = \lim_{\varepsilon \rightarrow 0} \delta_{\varepsilon^{-1}}^x \Delta^x(c(t), c(t + \varepsilon))$$

We use now the properties of Q^x in the left hand side of the previous equality:

$$Q^x \delta_{\varepsilon^{-1}}^x \Delta^x(c(t), c(t + \varepsilon)) = \bar{\delta}_{\varepsilon^{-1}}^x Q^x \Delta^x(c(t), c(t + \varepsilon)) = \bar{\delta}_{\varepsilon^{-1}}^x \bar{\Delta}^x(Q^x c(t), Q^x c(t + \varepsilon))$$

thus after taking the limit as $\varepsilon \rightarrow 0$ we prove that the limit $\lim_{\varepsilon \rightarrow 0} \bar{\delta}_{\varepsilon^{-1}}^x \bar{\Delta}^x(Q^x c(t), Q^x c(t + \varepsilon))$ exists and we obtain:

$$\lim_{\varepsilon \rightarrow 0} \delta_{\varepsilon^{-1}}^x \Delta^x(c(t), c(t + \varepsilon)) = \lim_{\varepsilon \rightarrow 0} \bar{\delta}_{\varepsilon^{-1}}^x \bar{\Delta}^x(Q^x c(t), Q^x c(t + \varepsilon))$$

This last equality is the same as (47), which is equivalent with (46). As a consequence we obtain the following equality, for almost any $t \in [0, 1]$:

$$\bar{d}^x \left(x, \Delta^x(c(t), \frac{\dot{d}^x}{dt} c(t)) \right) = \bar{\Delta}^x(Q^x c(t), \frac{\bar{d}^x}{dt} (Q^x c)(t)) \quad (48)$$

This implies that $Q^x c$ is absolutely continuous and by theorem 2.15, as in the proof of theorem 8.3 (but without using the Radon-Nikodym property, because we already know that $Q^x c$ is derivable a.e.), we obtain the following formula for the length of the curve $Q^x c$:

$$\bar{l}^x(Q^x c) = \int_0^1 \bar{d}^x \left(x, \bar{\Delta}^x(Q^x c(t), \frac{\bar{d}^x}{dt} (Q^x c)(t)) \right) dt$$

But we have also:

$$l^x(c) = \int_0^1 \bar{d}^x \left(x, \Delta^x(c(t), \frac{\dot{d}^x}{dt} c(t)) \right) dt$$

By (48) we obtain $\bar{l}^x(Q^x c) = l^x(c)$. \square

Proposition 12.7 *If $(X, \bar{d}, \bar{\delta})$ is a strong dilation structure, Q is a coherent projection and \bar{d}^x is finite then the triple $(U(x), \Sigma^x, \delta^x)$ is a normed conical group, with the norm induced by the left-invariant distance \bar{d}^x .*

Proof. The fact that $(U(x), \Sigma^x, \delta^x)$ is a conical group comes directly from the definition 10.2 of a coherent projection. Indeed, it is enough to use proposition 10.3 (c) and the formalism of binary decorated trees in [4] section 4 (or theorem 11 [4]), in order to reproduce the part of the proof of theorem 10 (p.87-88) in that paper, concerning the conical group structure. There is one small subtlety though. In the proof of theorem 7.3(a) the same modification of proof has been done starting from the axiom A4+, namely the existence of the uniform limit $\lim_{\varepsilon \rightarrow 0} \Sigma_{\varepsilon}^x(u, v) = \Sigma^x(u, v)$. Here we need first to prove this limit, in a similar way as in the corollary 9 [4]. We shall use for this the distance \bar{d}^x instead of the distance in the metric tangent space of (X, d) at x denoted by d^x (which is not yet proven to exist). The distance \bar{d}^x is supposed to be finite by hypothesis. Moreover, by its definition and theorem 12.6 we have

$$\bar{d}^x(u, v) \geq d^x(u, v)$$

therefore the distance \bar{d}^x is non degenerate. By construction this distance is also left invariant with respect to the group operation $\Sigma^x(\cdot, \cdot)$. Therefore we may repeat the proof of corollary 9 [4] and obtain the result that A4+ is true for (X, d, δ) .

What we need to prove next is that \mathring{d}^x induces a norm on the conical group $(U(x), \Sigma^x, \delta^x)$. For this it is enough to prove that

$$\mathring{d}^x(\mathring{\delta}_\mu^{x,u}v, \mathring{\delta}_\mu^{x,u}w) = \mu \mathring{d}^x(v, w) \quad (49)$$

for any $v, w \in U(x)$. This is a direct consequence of relation (48) from the proof of the theorem 12.6. Indeed, by direct computation we get that for any curve c which is \mathring{Q}^x -horizontal a.e. we have:

$$\begin{aligned} l^x(\mathring{\delta}_\mu^{x,u}c) &= \int_0^1 \bar{d}^x \left(x, \Delta^x \left(\mathring{\delta}_\mu^{x,u}c(t), \frac{\mathring{d}^x}{dt} \left(\mathring{\delta}_\mu^{x,u}c \right) (t) \right) \right) dt = \\ &= \int_0^1 \bar{d}^x \left(x, \delta_\mu^x \Delta^x \left(c(t), \frac{\mathring{d}^x}{dt} c(t) \right) \right) dt \end{aligned}$$

But c is \mathring{Q}^x -horizontal a.e., which implies, via (48), that

$$\delta_\mu^x \Delta^x \left(c(t), \frac{\mathring{d}^x}{dt} c(t) \right) = \bar{\delta}_\mu^x \Delta^x \left(c(t), \frac{\mathring{d}^x}{dt} c(t) \right)$$

therefore we have

$$l^x(\mathring{\delta}_\mu^{x,u}c) = \int_0^1 \bar{d}^x \left(x, \bar{\delta}_\mu^x \Delta^x \left(c(t), \frac{\mathring{d}^x}{dt} c(t) \right) \right) dt = \mu l^x(c)$$

This implies (49), therefore the proof is done. \square

Theorem 12.8 *If the generalized Chow condition (Cgen) and condition (B) are true then $(U(x), \Sigma^x, \delta^x)$ is a local conical group which is a neighbourhood of the neutral element of a Carnot group generated by $Q^x U(x)$.*

Proof. For any for any $\varepsilon \in (0, 1]$, $x, u, v \in X$ sufficiently close and $\mu > 0$ we have the relations:

$$(i) \quad \hat{Q}_{\varepsilon, \mu}^{x,u}v = \Sigma_\varepsilon^x(u, Q_\mu^{\delta_\varepsilon^{x,u}} \Delta_\varepsilon^x(u, v)),$$

$$(ii) \quad \hat{Q}_\varepsilon^{x,u}v = \Sigma_\varepsilon^x(u, Q^{\delta_\varepsilon^{x,u}} \Delta_\varepsilon^x(u, v)).$$

(i) implies (ii) when $\mu \rightarrow 0$, thus it is sufficient to prove only the first point. This is the result of a computation:

$$\begin{aligned} \hat{Q}_{\varepsilon, \mu}^{x,u}v &= \delta_{\varepsilon^{-1}}^x Q_\mu^{\delta_\varepsilon^{x,u}} \delta_\varepsilon^x = \\ &= \delta_{\varepsilon^{-1}}^x \delta_\varepsilon^{\delta_\varepsilon^{x,u}} Q_\mu^{\delta_\varepsilon^{x,u}} \delta_{\varepsilon^{-1}}^{\delta_\varepsilon^{x,u}} \delta_\varepsilon^x = \Sigma_\varepsilon^x(u, Q_\mu^{\delta_\varepsilon^{x,u}} \Delta_\varepsilon^x(u, v)) \end{aligned}$$

It follows that we can put the recurrence relations (44) in the form:

$$\Psi_{\varepsilon w}^{k+1}([q]_{k+1}) = \Sigma_\varepsilon^x \left(\Psi_{\varepsilon w}^k([q]_k), Q_{w_k}^{\delta_\varepsilon^{x, \Psi_{\varepsilon w}^k([q]_k)}} \Delta_\varepsilon^x \left(\Psi_{\varepsilon w}^k([q]_k), q_{k+1} \right) \right) \quad (50)$$

This recurrence relation allows us to prove by induction that for any k the limit

$$\Psi_w^k([q]_k) = \lim_{\varepsilon \rightarrow 0} \Psi_{\varepsilon w}^k([q]_k)$$

exists and it satisfies the recurrence relation:

$$\Psi_{0w}^{k+1}([q]_{k+1}) = \Sigma^x \left(\Psi_{0w}^k([q]_k), Q_{w_k}^x \Delta^x \left(\Psi_{0w}^k([q]_k), q_{k+1} \right) \right) \quad (51)$$

and the initial condition $\Psi_{0w}^1(x) = x$. We pass to the limit in the generalized Chow condition (Cgen) and we thus obtain that a neighbourhood of the neutral element x is (algebraically) generated by $Q^x U(x)$. Then the distance \bar{d}^x . Therefore by proposition 12.7 $(U(x), \Sigma^x, \delta^x)$ is a normed conical group generated by $Q^x U(x)$.

Let $c : [0, 1] \rightarrow U(x)$ be the curve $c(t) = \delta_t^x u$, with $u \in Q^x U(x)$. Then we have $Q^x c(t) = c(t) = \bar{\delta}_t^x u$. From condition (B) we get that c is $\bar{\delta}$ -derivable at $t = 0$. A short computation of this derivative shows that:

$$\frac{d\bar{\delta}}{dt} c(0) = u$$

Another easy computation shows that the curve c is $\bar{\delta}^x$ -derivable if and only if the curve c is $\bar{\delta}$ -derivable at $t = 0$, which is true, therefore c is $\bar{\delta}^x$ -derivable, in particular at $t = 0$. Moreover, the expression of the $\bar{\delta}^x$ -derivative of c shows that c is also Q^x -everywhere horizontal (compare with theorem 11.1). We use the theorem 12.6 and relation (46) from its proof to deduce that $c = Q^x c$ is δ^x -derivable at $t = 0$, thus for any $u \in Q^x U(x)$ and small enough $t, \tau \in (0, 1)$ we have

$$\delta_{t+\tau}^{\delta^x, x} u = \bar{\Sigma}^x(\bar{\delta}_t^x u, \bar{\delta}_\tau^x u) \quad (52)$$

By the previous proposition 12.7 and corollary 6.3 [5], the normed conical group $(U(x), \Sigma^x, \delta^x)$ is in fact locally a homogeneous group, i.e. a simply connected Lie group which admits a positive graduation given by the eigenspaces of δ^x . Indeed, corollary 6.3 [4] is originally about strong dilation structures, but the generalized Chow condition implies that the distances d , \bar{d} and \bar{d}^x induce the same uniformity, which, along with proposition 12.7, are the only things needed for the proof of this corollary. The conclusion of corollary 6.3 [5] therefore is true, that is $(U(x), \Sigma^x, \delta^x)$ is locally a homogeneous group. Moreover it is locally Carnot if and only if on the generating space $Q^x U(x)$ any dilation $\delta_\varepsilon^{\delta^x, x} u = \bar{\delta}_\varepsilon^x$ is linear in ε . But this is true, as shown by relation (52). This ends the proof. \square

12.3 Coherent projections induce length dilation structures

Theorem 12.9 *If $(X, \bar{d}, \bar{\delta})$ is a tempered strong dilation structure, has the Radon-Nikodym property and Q is a coherent projection, which satisfies (A), (B), (Cgen) then (X, d, δ) is a length dilation structure.*

Proof. We shall prove that:

- (a) for any function $\varepsilon \in (0, 1) \mapsto (x_\varepsilon, c_\varepsilon) \in \mathcal{L}_\varepsilon(X, d, \delta)$ which converges to (x, c) as $\varepsilon \rightarrow 0$, with $c : [0, 1] \rightarrow U(x)$ δ^x -derivable and \bar{Q}^x -horizontal almost everywhere, we have:

$$l^x(c) \leq \liminf_{\varepsilon \rightarrow 0} l^{x_\varepsilon}(c_\varepsilon)$$

- (b) for any sequence $\varepsilon_n \rightarrow 0$ and any (x, c) , with $c : [0, 1] \rightarrow U(x)$ δ^x -derivable and \bar{Q}^x -horizontal almost everywhere, there is a recovery sequence $(x_n, c_n) \in \mathcal{L}_{\varepsilon_n}(X, d, \delta)$ such that

$$l^x(c) = \lim_{n \rightarrow \infty} l^{x_n}(c_n)$$

Proof of (a). This is a consequence of theorem 12.6 and definition 10.2 of a coherent projection. With the notations from (a), let us first prove that $l^x(c) = \bar{l}^x(Q^x c)$. Let c be a curve such that $\delta_\varepsilon^x c$ is \bar{d} -Lipschitz and Q -horizontal. Then:

$$l_\varepsilon^x(c) = \sup \left\{ \sum_{i=1}^n \frac{1}{\varepsilon} \bar{d}(\delta_\varepsilon^x c(t_i), \delta_\varepsilon^x c(t_{i+1})) : 0 = t_1 < \dots < t_{n+1} = 1 \right\} =$$

$$= \sup \left\{ \sum_{i=1}^n \frac{1}{\varepsilon} \bar{d}(\bar{\delta}_\varepsilon^x Q_\varepsilon^x c(t_i), \bar{\delta}_\varepsilon^x Q_\varepsilon^x c(t_{i+1})) : 0 = t_1 < \dots < t_{n+1} = 1 \right\} = \\ = \bar{l}_\varepsilon^x(Q_\varepsilon^x c)$$

Now we have to prove the following:

$$l^x(c) = \bar{l}^x(Q^x c) \leq \liminf_{\varepsilon \rightarrow 0} \bar{l}^{x_\varepsilon}(Q_\varepsilon^{x_\varepsilon} c_\varepsilon)$$

This is true because $(X, \bar{d}, \bar{\delta})$ is a tempered dilation structure and because of condition (A). Indeed from the fact that $(X, \bar{d}, \bar{\delta})$ is tempered and from (41) (which is a consequence of condition (A)) we deduce that Q_ε is uniformly continuous on compact sets in a uniform way: for any compact set $K \subset X$ there are constants $L(K) > 0$ (from (A)) and $C > 0$ (from the tempered condition) such that for any $\varepsilon \in (0, 1]$, any $x \in K$ and any u, v sufficiently close to x we have:

$$\bar{d}(Q_\varepsilon^x u, Q_\varepsilon^x v) \leq C (\bar{\delta}_\varepsilon^x \bar{d})(Q_\varepsilon^x u, Q_\varepsilon^x v) \leq C L(K) \bar{d}(u, v)$$

The sequence Q_ε^x uniformly converges to Q^x as ε goes to 0, uniformly with respect to x in compact sets. Therefore if $(x_\varepsilon, c_\varepsilon) \in \mathcal{L}_\varepsilon(X, d, \delta)$ converges to (x, c) then $(x_\varepsilon, Q_\varepsilon^{x_\varepsilon} c_\varepsilon) \in \mathcal{L}_\varepsilon(X, \bar{d}, \bar{\delta})$ converges to $(x, Q^x c)$. Use now the fact that by corollary 8.9 $(X, \bar{d}, \bar{\delta})$ is a length dilation structure. The proof is done.

Proof of (b). We have to construct a recovery sequence. We are doing this by discretization of $c : [0, L] \rightarrow U(x)$. Recall that c is a curve which is δ^x -derivable a.e. and \dot{Q}^x -horizontal, that is for almost every $t \in [0, L]$ the limit

$$u(t) = \lim_{\mu \rightarrow 0} \delta_{\mu^{-1}}^x \Delta^x(c(t), c(t + \mu))$$

exists and $Q^x u(t) = u(t)$. Moreover we may suppose that for almost every t we have $\bar{d}^x(x, u(t)) \leq 1$ and $\bar{l}^x(c) \leq L$.

There are functions $\omega^1, \omega^2 : (0, +\infty) \rightarrow [0, +\infty)$ with $\lim_{\lambda \rightarrow 0} \omega^i(\lambda) = 0$, with the following property: for any $\lambda > 0$ sufficiently small there is a division $A_\lambda = \{0 < t_0 < \dots < t_P < L\}$ such that

$$\frac{\lambda}{2} \leq \min \left\{ \frac{t_0}{t_1 - t_0}, \frac{L - t_P}{t_P - t_{P-1}}, t_k - t_{k-1} : k = 1, \dots, P \right\} \quad (53)$$

$$\lambda \geq \max \left\{ \frac{t_0}{t_1 - t_0}, \frac{L - t_P}{t_P - t_{P-1}}, t_k - t_{k-1} : k = 1, \dots, P \right\} \quad (54)$$

and such that $u(t_k)$ exists for any $k = 1, \dots, P$ and

$$\dot{d}^x(c(0), c(t_0)) \leq t_0 \leq \lambda^2 \quad (55)$$

$$\dot{d}^x(c(L), c(t_P)) \leq L - t_P \leq \lambda^2 \quad (56)$$

$$\dot{d}^x(u(t_{k-1}), \Delta^x(c(t_{k-1}), c(t_k))) \leq (t_k - t_{k-1}) \omega^1(\lambda) \quad (57)$$

$$\left| \int_0^L \bar{d}^x(x, u(t)) dt - \sum_{k=0}^{P-1} (t_{k+1} - t_k) \bar{d}^x(x, u(t_k)) \right| \leq \omega^2(\lambda) \quad (58)$$

Indeed (55), (56) are a consequence of the fact that c is \dot{d}^x -Lipschitz, (57) is a consequence of Egorov theorem applied to

$$f_\mu(t) = \delta_{\mu^{-1}}^x \Delta^x(c(t), c(t + \mu))$$

and (58) comes from the definition of the integral

$$l(c) = \int_0^L \bar{d}^x(x, u(t)) dt$$

For each λ we shall choose $\varepsilon = \varepsilon(\lambda)$ and we shall construct a curve c_λ with the properties:

- (i) $(x, c_\lambda) \in \mathcal{L}_{\varepsilon(\lambda)}(X, d, \delta)$
- (ii) $\lim_{\lambda \rightarrow 0} l_{\varepsilon(\lambda)}^x(c_\lambda) = l^x(c)$.

At almost every t the point $u(t)$ represents the velocity of the curve c seen as the the left translation of $\frac{d^x}{dt}c(t)$ by the group operation $\Sigma^x(\cdot, \cdot)$ to x (which is the neutral element for the mentioned operation). The derivative (with respect to δ^x) of the curve c at t is

$$y(t) = \Sigma^x(c(t), u(t))$$

Let us take $\varepsilon > 0$, arbitrary for the moment. We shall use the points of the division A_λ and for any $k = 0, \dots, P - 1$ we shall define the point:

$$y_k^\varepsilon = \hat{Q}_\varepsilon^{x, c(t_k)} \Sigma_\varepsilon^x(c(t_k), u(t_k)) \quad (59)$$

Thus y_k^ε is obtained as the "projection" by $\hat{Q}_\varepsilon^{x, c(t_k)}$ of the "approximate left translation" $\Sigma_\varepsilon^x(c(t_k), \cdot)$ by $c(t_k)$ of the velocity $u(t_k)$. Define also the point:

$$y_k = \Sigma^x(c(t_k), u(t_k))$$

By construction we have:

$$y_k^\varepsilon = \hat{Q}_\varepsilon^{x, c(t_k)} y_k \quad (60)$$

and by computation we see that y_k^ε can be expressed as:

$$\begin{aligned} y_k^\varepsilon &= \delta_{\varepsilon^{-1}}^x Q^{\delta_\varepsilon^x c(t_k)} \delta_{\varepsilon^{\delta_\varepsilon^x c(t_k)}}^{\delta_\varepsilon^x} u(t_k) = \\ &= \Sigma_\varepsilon^x(c(t_k), Q^{\delta_\varepsilon^x c(t_k)} u(t_k)) = \delta_{\varepsilon^{-1}}^x \bar{\delta}_{\varepsilon^{\delta_\varepsilon^x c(t_k)}}^{\delta_\varepsilon^x} Q^{\delta_\varepsilon^x c(t_k)} u(t_k) \end{aligned} \quad (61)$$

Let us define the curve

$$c_k^\varepsilon(s) = \hat{\delta}_{\varepsilon, s}^{x, c(t_k)} y_k^\varepsilon \quad , \quad s \in [0, t_{k+1} - t_k] \quad (62)$$

which is a \hat{Q}_ε^x -horizontal curve (by supplementary hypothesis (B)) which joins $c(t_k)$ with the point

$$z_k^\varepsilon = \hat{\delta}_{\varepsilon, t_{k+1} - t_k}^{x, c(t_k)} y_k^\varepsilon \quad (63)$$

The point z_k^ε is an approximation of the point

$$z_k = \hat{\delta}_{t_{k+1} - t_k}^{x, c(t_k)} y_k$$

We shall also consider the curve

$$c_k(s) = \hat{\delta}_s^{x, c(t_k)} y_k \quad , \quad s \in [0, t_{k+1} - t_k] \quad (64)$$

There is a short curve g_k^ε which joins z_k^ε with $c(t_{k+1})$, according to condition (Cgen). Indeed, for ε sufficiently small the points $\delta_\varepsilon^x z_k^\varepsilon$ and $\delta_\varepsilon^x c(t_{k+1})$ are sufficiently close.

Finally, take g_0^ε and g_{P+1}^ε "short curves" which join $c(0)$ with $c(t_0)$ and $c(t_P)$ with $c(L)$ respectively.

Correspondingly, we can find short curves g_k (in the geometry of the dilation structure $(U(x), \bar{d}^x, \hat{\delta}^x, \hat{Q}^x)$) joining z_k with $c(t_{k+1})$, which are the uniform limit of the short curves g_k^ε as $\varepsilon \rightarrow 0$. Moreover this convergence is uniform with respect to k (and λ). Indeed, these short curves are made by N curves of the type $s \mapsto \hat{\delta}_{\varepsilon, s}^{x, u_\varepsilon} v_\varepsilon$, with $\hat{Q}^{x, u_\varepsilon} v_\varepsilon = v_\varepsilon$. Also, the short curves g_k are made respectively by N curves of the type $s \mapsto \hat{\delta}_s^{x, u} v$, with $\hat{Q}^{x, u} v = v$. Therefore we have:

$$\begin{aligned} & \bar{d}(\hat{\delta}_s^{x, u} v, \hat{\delta}_{\varepsilon, s}^{x, u_\varepsilon} y_k^\varepsilon) = \\ & = \bar{d}(\Sigma^x(u, \bar{\delta}_s^x \Delta^x(u, v)), \Sigma_\varepsilon^x(u_\varepsilon, \bar{\delta}_{s^\varepsilon}^{x, u_\varepsilon} \Delta_\varepsilon^x(u_\varepsilon, v_\varepsilon))) \end{aligned}$$

By an induction argument on the respective ends of segments forming the short curves, using the axioms of coherent projections, we get the result.

By concatenation of all these curves we get two new curves:

$$\begin{aligned} c_\lambda^\varepsilon &= g_0^\varepsilon \left(\prod_{k=0}^{P-1} c_k^\varepsilon g_k^\varepsilon \right) g_{P+1}^\varepsilon \\ c_\lambda &= g_0 \left(\prod_{k=0}^{P-1} c_k g_k \right) g_{P+1} \end{aligned}$$

From the previous reasoning we get that as $\varepsilon \rightarrow 0$ the curve c_λ^ε uniformly converges to c_λ , uniformly with respect to λ .

By theorem 12.8, specifically from relation (52) and considerations below, we notice that for any $u = Q^x u$ the length of the curve $s \mapsto \delta_s^x u$ is:

$$l^x(s \in [0, a] \mapsto \delta_s^x u) = a \bar{d}^x(x, u)$$

From here and relations (55), (56), (57), (58) we get that

$$l^x(c) = \lim_{\lambda \rightarrow 0} l^x(c_\lambda) \quad (65)$$

Condition (B) and the fact that $(X, \bar{d}, \bar{\delta})$ is tempered imply that there is a positive function $\omega^3(\varepsilon) = \mathcal{O}(\varepsilon)$ such that

$$|l_\varepsilon^x(c_\lambda^\varepsilon) - l^x(c_\lambda)| \leq \frac{\omega^3(\varepsilon)}{\lambda} \quad (66)$$

This is true because if $v \hat{Q}_\varepsilon^{x, u} v$ then $\delta_\varepsilon^x v = Q^{\delta_\varepsilon^x u} \delta_\varepsilon^x v$, therefore by condition (B)

$$\frac{l_\varepsilon^x(s \in [0, a] \mapsto \hat{\delta}_{\varepsilon, s}^{x, u} v)}{\delta_\varepsilon^x \bar{d}(u, v)} = \frac{\bar{l}(s \in [0, a] \mapsto \bar{\delta}_s^{\delta_\varepsilon^x u} \delta_\varepsilon^x v)}{\bar{d}(\delta_\varepsilon^x u, \delta_\varepsilon^x v)} \leq \mathcal{O}(\varepsilon) + 1$$

Since each short curve is made by N segments and the division A_λ is made by $1/\lambda$ segments, the relation (66) follows.

We shall choose now $\varepsilon(\lambda)$ such that $\omega^3(\varepsilon(\lambda)) \leq \lambda^2$ and we define:

$$c_\lambda = c_\lambda^{\varepsilon(\lambda)}$$

These curves satisfy the properties (i), (ii). Indeed (i) is satisfied by construction and (ii) follows from the choice of $\varepsilon(\lambda)$, uniform convergence of c_λ^ε to c_λ , uniformly with respect to λ , and relations (66), and (65). \square

References

- [1] A.D. Alexandrov, Intrinsic geometry of convex surfaces, Akademie Verlag, Berlin (1955)
- [2] L. Ambrosio, N. Gigli, G. Savaré, Gradient flows in metric spaces and in the space of probability measures, Birkhäuser Verlag, Basel-Boston-Berlin, (2005)
- [3] A. Bellaïche, The tangent space in sub-Riemannian geometry, in: Sub-Riemannian Geometry, A. Bellaïche, J.-J. Risler eds., *Progress in Mathematics*, **144**, Birkhäuser, (1996), 4-78
- [4] M. Buliga, Dilatation structures I. Fundamentals, *J. Gen. Lie Theory Appl.*, Vol **1** (2007), No. 2, 65-95. <http://arxiv.org/abs/math.MG/0608536>
- [5] M. Buliga, Infinitesimal affine geometry of metric spaces endowed with a dilation structure (2008), *Houston Journal of Math.*, **36**, 1 (2010), 91-136, <http://arxiv.org/abs/0804.0135>
- [6] M. Buliga, Dilatation structures in sub-riemannian geometry, (2007), in: Contemporary Geometry and Topology and Related Topics, Cluj-Napoca, Cluj University Press (2008), 89-105 , <http://arxiv.org/abs/0708.4298>
- [7] M. Buliga, A characterization of sub-riemannian spaces as length dilation structures constructed via coherent projections, *Commun. Math. Anal.*, **11** (2011), No. 2, pp. 70-111
- [8] M. Buliga, Braided spaces with dilations and sub-riemannian symmetric spaces. in: Geometry. Exploratory Workshop on Differential Geometry and its Applications, eds. D. Andrica, S. Moroianu, Cluj-Napoca 2011, 21-35, <http://arxiv.org/abs/1005.5031>
- [9] M. Buliga, Self-similar dilatation structures and automata, Proceedings of the 6-th Congress of Romanian Mathematicians, Bucharest, 2007, vol. 1, 557-564, <http://fr.arxiv.org/abs/0709.2224> (2007)
- [10] M. Buliga, Tangent bundles to sub-Riemannian groups (2003), <http://arxiv.org/abs/math/0307342>
- [11] M. Buliga, Curvature of sub-Riemannian spaces (2003), <http://arxiv.org/abs/math/0311482>
- [12] M. Buliga, Sub-Riemannian geometry and Lie groups. Part I. (2002), <http://arxiv.org/abs/math/0210189>
- [13] M. Buliga, Sub-Riemannian geometry and Lie groups. Part II. Curvature of metric spaces, coadjoint orbits and associated representations, (2004), <http://arxiv.org/abs/math/0407099>
- [14] M. Buliga, Normed groupoids with dilations, (2011), <http://arxiv.org/abs/1107.2823>
- [15] M. Buliga, Maps of metric spaces, (2011), <http://arxiv.org/abs/1107.2817>
- [16] G. Buttazzo, L. De Pascale, I. Fragalà, Topological equivalence of some variational problems involving distances, *Discrete Contin. Dynam. Systems* **7** (2001), no. 2, 247-258
- [17] W.L. Chow, Über Systeme von linearen partiellen Differentialgleichungen erster Ordnung, *Math. Ann.*, **117** (1939), 98-105.
- [18] G. Dal Maso, An introduction to Γ -convergence. Progress in Nonlinear Differential Equations and Their Applications **8**, Birkhäuser, Basel (1993)

- [19] G. David, S. Semmes, Fractured fractals and broken dreams: Self-similar geometry through metric and measure, Clarendon Press, (Oxford and New York) 1997
- [20] L. van den Dries, I. Goldbring, Locally compact contractive local groups, *J. of Lie Theory*, **19** (2009), 685-695, <http://arxiv.org/abs/0909.4565>
- [21] G.B. Folland, E.M. Stein, Hardy spaces on homogeneous groups, Mathematical Notes, **28**, Princeton University Press, N.J.; University of Tokyo Press, Tokyo, 1982.
- [22] M. Fréchet, Sur quelques points du calcul fonctionnel, *Rendic. Circ. Mat. Palermo* **22** (1906), 1-72
- [23] H. Glöckner, Contractible Lie groups over local fields, (2007), available as e-print <http://arxiv.org/abs/0704.3737>
- [24] H. Glöckner, G.A. Willis, Classification of the simple factors appearing in composition series of totally disconnected contraction groups, (2006), available as e-print <http://arxiv.org/abs/math/0604062>
- [25] M. Gromov, Carnot-Carathéodory spaces seen from within, in the book: Sub-Riemannian Geometry, A. Bellaïche, J.-J. Risler eds., *Progress in Mathematics*, **144**, Birkhäuser, (1996), 79-323.
- [26] M. Gromov, Metric structures for Riemannian and non-Riemannian spaces, Progress in Math., **152**, Birkhäuser (1999)
- [27] J. Mitchell, On Carnot-Carathéodory metrics, *Journal of Differential Geom.*, **21** (1985), 35-45.
- [28] I.G. Nikolaev, A metric characterization of riemannian spaces, *Siberian Adv. Math.* , **9** (1999), 1-58
- [29] P. Pansu, Métriques de Carnot-Carathéodory et quasi-isométries des espaces symétriques de rang un, *Ann. of Math.*, (2) **129**, (1989), 1-60
- [30] E. Siebert, Contractive automorphisms on locally compact groups, *Math. Z.*, **191**, 73-90, (1986)
- [31] S. Venturini, Derivation of distance functions in \mathbb{R}^n , preprint (1991)
- [32] S.K. Vodopyanov, Differentiability of mappings in the geometry of the Carnot manifolds, *Siberian Math. J.*, Vol. **48**, No. 2, (2007), 197-213
- [33] S.K. Vodopyanov, M. Karmanova, Local geometry of Carnot manifolds under minimal smoothness, *Doklady Math.* **412**, 3, 305-311, (2007)
- [34] A. Wald, Begründung einer koordinatenlosen Differentialgeometrie der Flächen, *Erg. Math. Colloq.*, **7** (1936), 24-46
- [35] J.S.P. Wang, The Mautner phenomenon for p-adic Lie groups, *Math. Z.* **185** (1966), 403-411