Traffic-related air pollution. A pilot exposure assessment in Beirut, Lebanon

Mireille Borgie, Anne Garat, Fabrice Cazier, Agnes Delbende, Delphine Allorge, Frederic Ledoux, Dominique Courcot, Pirouz Shirali, Zeina Dagher

► To cite this version:

Mireille Borgie, Anne Garat, Fabrice Cazier, Agnes Delbende, Delphine Allorge, et al.. Traffic-related air pollution. A pilot exposure assessment in Beirut, Lebanon. Chemosphere, 2013, pp.122-128. hal-00914713

HAL Id: hal-00914713

<https://hal-confremo.archives-ouvertes.fr/hal-00914713>

Submitted on 25 Jan 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

TRAFFIC-RELATED AIR POLLUTION. A PILOT EXPOSURE

ASSESSMENT IN BEIRUT, LEBANON

**Mireille BORGIE^{a,d}, Anne GARAT^{c,d}, Fabrice CAZIER^{b,d}, Agnes DELBENDE^{b,d},
Camille RICHEVAL^{c,d}, Delphine ALLORGE^{c,d}, Frederic LEDOUX^{a,d}, Dominique
COURCOT^{a,d}, Pirouz SHIRALI^{a,d}, Zeina DAGHER^{e,f}**

^a Unit of Environmental Chemistry and Interactions on Living, EA 4492, University of
Littoral-Côte d'Opale (ULCO), Dunkerque, France

^b Common Center of Measurements, ULCO, Dunkerque, France

^c EA 4483, Faculty of Medicine of Lille, University of Lille North of France, Lille, France

^d University of Lille North of France, Lille, France

^e Bioactive Molecules Research Group, Doctoral School of Sciences and Technologies,
Lebanese University, Lebanon

^f Department of Biology, Faculty of Sciences, Lebanese University, Lebanon

Corresponding Author: Dr Zeina DAGHER, Bioactive Molecules Research Group,
Doctoral School of Sciences and Technologies; Department of Biology, Faculty of Sciences,
Lebanese University, P.O. Box 90656, Lebanon. Tel +961 (1) 686 981, fax +961 (1) 680 250,
email: dagher_z@ul.edu.lb

23 ABSTRACT

24 Traffic-related volatile organic compounds (VOCs) pollution has frequently been
25 demonstrated to be a serious problem in the developing countries. Benzene and 1,3-butadiene
26 (BD) have been classified as a human carcinogen based on evidence for an increased
27 genotoxic and epigenotoxic effects in both occupational exposure assessment and *in vivo/in*
28 *vitro* studies. We have undertaken a biomonitoring of 25 traffic policemen and 23 office
29 policemen in Beirut, through personal air monitoring, assessed by diffusive samplers, as well
30 as through the use of biomarkers of exposure to benzene and BD. Personal benzene, toluene,
31 ethylbenzene, and xylene (BTEX) exposure were quantified by GC-MS/MS, urinary trans,
32 trans-muconic acid (*t,t*-MA) by HPLC/UV, S-phenyl mercapturic acid (S-PMA),
33 monohydroxy-butenyl mercapturic acid (MHBMA) and dihydroxybutyl mercapturic acid
34 (DHBMA) by ultra-performance liquid chromatography-electrospray tandem mass
35 spectrometry (UPLC/ESI(-)-MS/MS) in MRM (Multiple Reaction Monitoring) mode. We
36 found that individual exposure to benzene in the traffic policemen was higher than that
37 measured in traffic policemen in Prague, in Bologna, in Ioannina and in Bangkok. *t,t*-MA
38 levels could distinguish between office and traffic policemen and showed a better correlation
39 with personal BTEX compounds exposure. However, median MHBMA levels in traffic
40 policemen were slightly elevated, though not significantly higher than in office policemen.
41 Alternatively, DHBMA concentrations could significantly distinguish between office and
42 traffic policemen and showed a better correlation with personal total BTEX exposure.
43 DHMBA, measured in the post-shift urine samples, correlated with both pre-shift MHMBA
44 and pre-shift DHMBA. Moreover, there was not a marked effect of smoking habits on
45 DHBMA. Taken together, these findings suggested that DHBMA is more suitable than
46 MHBMA as biomarker of exposure to BD in humans. Traffic policemen, who are exposed to

47 benzene and BD at the roadside in central Beirut, are potentially at a higher risk for
48 development of diseases such as cancer than office policemen.

49

50 **Keywords:** Biomarkers; benzene; 1,3-butadiene; occupational exposure; traffic-related air
51 pollution

52

1. INTRODUCTION

Traffic-related volatile organic compounds (VOCs) pollution has frequently been demonstrated to be a more serious problem in the developing countries than in the United States and Europe, as indicated by the VOC data obtained in Thailand, India, Pakistan and Egypt (Arayasiri et al., 2010; Rekhadevi et al., 2010; Kamal et al., 2012; Ibrahim et al., 2012). In Beirut, capital of Lebanon, air pollutant concentrations currently exceed air quality standards and guidelines (Waked and Afif, 2012). About 67% of non methanic VOC emissions are calculated to originate from the on-road transport sector and the majority of vehicles operate on gasoline (Waked and Afif, 2012).

Since concentrations of VOCs are elevated, albeit to different extents, on and near roadways, the individuals whose job requires that they spend long periods of time near vehicles may incur substantial occupational exposures to traffic-related air pollution (Knibbs and Morawska et al., 2012). It is well known that exposure data from stationary monitoring sites cannot give the real exposure profile in urban areas, since the level of traffic VOCs decreases drastically as the distance from the main traffic roads increases (Han and Naeher, 2006). More and until now, no studies have been conducted to assess of the human health risks from urban air pollution exposure in Beirut. Taken together, we carried out a personal exposure measurement campaign among traffic policemen to benzene and 1,3-butadiene (BD), since both compounds are generated from the incomplete combustion of gasoline.

Benzene and BD have been classified as Group 1 carcinogens (IARC; 2008, 2009) based on evidence for an increased genotoxic and epigenotoxic effects in both occupational exposure assessment (Ruchirawat et al., 2010; Carugno et al., 2012; Peluso et al., 2012; Seow et al., 2012; Xiang et al., 2012) and in *in vivo* and *in vitro* studies (Dagher et al., 2006; Billet et al., 2010; Koturbash et al., 2011; Sangaraju et al., 2012; Tabish et al., 2012; Abbas et al., 2013).

Biomarkers of benzene, as urinary trans, trans-muconic acid (*t,t*-MA) and urinary S-phenylmercapturic acid (S-PMA), have been measured mostly in fuel-related exposure such as in station attendants, in public transportation or in traffic policemen (Fustinoni et al., 2005; Barbieri et al., 2008; Manini et al., 2010), while only a few studies of traffic-related exposures to BD have been performed (Sapkota et al., 2006; Arayasiri et al., 2010).

Although, BD is a known human carcinogen emitted from mobile sources, little is known about traffic-related human exposure to this toxicant. BD is metabolized *in vivo* to reactive epoxides which are supposedly responsible for the observed carcinogenic effects (category 1A; EU-RAR 2002). A main metabolic pathway for these epoxides is the reaction with glutathione, leading to a urinary excretion of 3,4-dihydroxybutyl mercapturic acid and 3-monohydroxybutenyl mercapturic acids (DHBMA and MHBMA). Up to now, DHBMA and MHBMA have already been used in population surveys as a biomarker of exposure to BD (Arayasiri et al., 2010; Ruchirawat et al., 2010; Cheng et al., 2012).

Hence, it will be of great interest to conduct a biomonitoring pilot study in order to assess traffic-related VOCs in central Beirut, and to evaluate the use of biomarkers of benzene and BD exposure of urban traffic policemen. This work was, therefore, undertaken to determine *t,t*-MA, SPMA, MHBMA and DHBMA in urine spot samples before and at the end of a working shift, and personal air monitoring to airborne benzene, toluene, ethylbenzene, and xylene (BTEX) during the work shift. The influence of personal exposure, job activity and personal characteristics on biomarkers excretion was evaluated.

2. MATERIALS AND METHODS

2.1. Measurements campaign design

The campaign took place during May-June 2011 and included measurements of BTEX personal exposure to 47 healthy volunteers. All participants were males. The volunteers group includes 24 traffic policemen and 23 office policemen which constitute the control group. The traffic policemen group consisted of officers whose activity consists exclusively of traffic regulation at intersections of central roads in the city. All participants were carrying passive samplers for BTEX during the working hours.

All participants kept a questionnaire requesting information about lifestyle and health status and a personal daily questionnaire, where they referred to the duration of the performed activities during sampling time.

2.2. Sample collection

The sampling was conducted on Monday, which is the first day of the work week after a two days holiday over the weekend to minimize residual exposure from the previous week. The work shift was 7 h/d and 5 h/d for traffic and office policemen, respectively. Individual air samples were attached to the clothing in the breathing zone of study subjects and throughout the entire work shift. After air sampling was completed, samples were capped, transported to the laboratory, and stored at 4 °C until analysis within 8 days by the end of the sampling campaign. Urine samples were collected at both pre-shift and post-shift and stored at -80 °C until analysis. All participants gave their informed consent and the study was approved by the Ethics Committee of the Lebanese University.

2.3. Analysis of BTEX

BTEX compounds were analyzed as described previously by Avogbe and colleagues (2011). Exposure to airborne BTEX was assessed by using GABIE (Gas Adsorbent Badge for Individual Exposure) diffusive samplers (ARELCO, ARC20001UP, France) containing activated charcoal cartridge. After sampling, the badges were sealed, preserved at -80 °C and sent for analysis to the “Centre Commun de Mesure”, ULCO, Dunkerque (France). Briefly, BTEX were desorbed from the activated charcoal by using 2 mL of benzene-free carbon disulfide (Sigma, France) under agitation for 15 min. The mixture was filtered and 1 µL of the filtrate was analysed on a Gas Chromatograph (GC) (CP-3800, Varian USA) coupled to a Mass Spectrometer (1200 TQ, Varian USA) using Factor four VF-5 ms column (0.25 mm internal diameter, 30 m, film thickness 0.25 µm). The carrier gas was helium, and the flow rate was set at 1 mL/min. The GC oven was held at 40 °C for 5 min and then increased to 310 °C at the rate of 5 °C/min. The recovery rate of extraction of benzene was 99%. Data were averaged over each sampling period.

2.4. Determination of urinary metabolites

2.4.1. Urinary *t,t*-muconic acid (*t,t*-MA)

Urine samples were thawed at room temperature for 15 min with frequent stirring, and then centrifuged at 3000 g for 10 min. Aliquots of alkalinized urine were applied to a strong anion exchange (SAX 500 mg 3cc) column (Varian) and subsequently washed with 3 mL 1% (v/v) acetic acid. The *t,t*-muconic acid was eluted with 3 mL 10% (v/v) aqueous acetic acid and then analyzed by HPLC equipped with a UV detector (Waters, Milford, USA). The concentration of *t,t*-MA was expressed as µg/g creatinine. The limit of detection (LOD) was 3 µg L⁻¹. The limit of quantification (LOQ) was 10 µg L⁻¹. The coefficient of variation of the method was within 12% for inter-day determination.

2.4.2. Urinary S-phenylmercapturic acid (S-PMA)

Four hundred microlitres of urine containing 10 μL of deuterated internal standards at 10 $\mu\text{g mL}^{-1}$ were added to 0.400 mL of sodium acetate buffer at 10 mM and 1 mL of distilled water. Sample mixture was homogenized for 0.5 min and centrifuged at 5000 rpm for 5 min. The supernatant was loaded onto Oasis® MAX 3 cc/60 mg cartridge (Waters, Milford, USA) pre-conditioned with 2 mL of methanol (Chromanorm HPLC grade) followed by 2 mL of deionized water (Versol®, Aguetant, Lyon, France). The extraction cartridge was washed successively with 2 mL of distilled water and 2 mL of methanol. Following the final wash, the cartridges were dried for 1 min. Analytes elution was effected using 2 mL of a 1% formic acid in methanol (v/v). The extracted fraction was evaporated under a stream of nitrogen gas, then dissolved in 0.1 mL 0.1% formic acid/acetonitril (95:5 v/v) (Sigma-Aldrich, France).

Urinary S-PMA was analyzed using a Waters Acquity ultra-performance liquid chromatography (UPLC) performed on a Acquity UPLC BEH C18 (1.7 μm , 2.1 x 100 mm) at 50 °C. Samples (15 μL) were injected onto the column using a gradient elution of acetonitril and 0.01% aqueous acetic acid at a flow rate of 0.4 mL/min. Mass spectrometric selective detection was provided by a Waters Acquity Xevo TQD tandem mass spectrometer running in negative mode. Negative ions were acquired using electrospray ionization operated in the MRM (Multiple Reaction Monitoring) mode. The concentration of S-PMA was expressed as $\mu\text{g g}^{-1}$ creatinine. The LOQ was 1 $\mu\text{g L}^{-1}$.

2.4.3. Urinary monohydroxy-butenyl mercapturic acid (MHBMA) and dihydroxybutyl mercapturic acid (DHBMA)

The preparation of urine samples for analysis of MHBMA and DHBMA was carried out as previously described for the determination of urinary S-PMA. After the clean-up procedure, samples (15 μL) were injected onto the column using a gradient elution of

acetonitril and 0.1% aqueous formic acid at a flow rate of 0.4 mL/min and were subjected to analysis by a Waters Acquity TQD tandem mass spectrometer operating in ESI-MRM mode. Concentrations of urinary MHBMA and DHBMA were expressed as $\mu\text{g g}^{-1}$ creatinine and the LOQ was $5 \mu\text{g L}^{-1}$ and $50 \mu\text{g L}^{-1}$, respectively.

2.4.4. Urinary creatinine

The concentration of creatinine in the urine samples was performed by the Jaffé method using a Roche Diagnostics kit (Roche Diagnostics, France). The LOD was $34 \mu\text{g L}^{-1}$.

2.4.5. Cotinine assay

Urinary cotinine levels, as the major nicotine metabolite, were analyzed using DRI cotinine enzyme immunoassay (Microgenics GmbH, Passau, Germany) to check the tobacco smoke exposure reported in the lifestyle questionnaires. Subjects with cotinine levels greater than $500 \mu\text{g g}^{-1}$ of creatinine were considered active smokers. The LOD was $34 \mu\text{g L}^{-1}$. All the tests were performed according to the manufacturer's instructions.

2.5. Statistical Analysis

The Chi2 test was used for comparing categorical variables between groups; when expected values within cells are < 5 , Fisher exact test was used. For quantitative variables with normal distribution, Student test or ANOVA were used to compare between two groups. In case of non normal distribution, the Mann-Whitney or Kruskal-Wallis test were used to compare between mean ranks for two or different groups. Correlation analysis was performed with Spearman rank order correlation. Multivariate analysis was also performed to estimate the influence of BTEX levels on biomarkers. We adjusted our model on several possible potential confounders including, age, BMI, estimated vehicle counts for the exposure period,

199 smoking and alcohol consumption. A level of $p < 0.05$ was considered statistically significant
200 (SPSS version 16.0 software).
201

3. RESULTS

3.1. Demographic characteristics

Table 1 represents the distribution of subjects with respect to age, smoking, duration of exposure and body mass index (BMI). Statistical results for alcohol current use are not shown due to poor positive responses to the related question. The two groups studied had similar demographic characteristics. On an average, the work shift was 7 h/d and 5 h/d for traffic and office policemen, respectively. All participants were of the same social class. Within both groups, the length of employment was 4.47 ± 2.91 years (mean $\pm$ SD). Urinary cotinine levels were 890 ± 999 and $786 \pm 770 \mu\text{g g}^{-1}$ creatinine for smokers and 36 ± 15.5 and $27 \pm 6 \mu\text{g g}^{-1}$ creatinine for nonsmokers sampled from traffic and office policemen, respectively. In line with WHO (WHO, 1996) recommendations, only urine samples with creatinine concentration in the range 0.3–3.0 g L⁻¹ were considered.

3.2. BTEX in breathing zone air

The results of personal exposure monitoring are presented in Table 2. The levels of BTEX compounds exposure were much higher in traffic policemen than in the office population. Traffic policemen were exposed to significantly higher (14-fold) BTEX compounds concentrations than office policemen ($p < 0.001$). Traffic policemen were currently exposed to a wide range of benzene, toluene, ethylbenzene, m-xylene, o-xylene and p-xylene levels.

A close examination at the results reveals that individual exposure to benzene values are high to previously campaigns conducted in Bologna/Italy (Maffei et al., 2005), in Ioannina/Greece (Pilidis et al., 2008), in Prague/Czech Republic (Rossnerova et al., 2009) and in Bangkok/Thailand (Arayasiri et al., 2010). It is necessary to clarify that policemen

working outdoor were performing exclusively traffic regulation throughout the entire exposure period, during which the policemen stay in the middle of the intersections and they are exposed to the direct emissions of vehicles, where the concentrations are elevated. In addition, linear regression analysis estimated that each exposure to one vehicle increases the level of individual exposure to benzene, to o-xylene and to p-xylene of $3.23 \mu\text{g m}^{-3}$ ($p = 0.017$), $2.6 \mu\text{g m}^{-3}$ ($p = 0.05$) and $3.16 \mu\text{g m}^{-3}$ ($p = 0.05$), respectively. No significant relationships were observed between individual toluene or ethylbenzene exposure levels and estimated vehicle counts for the exposure period.

Moreover, excellent spearman correlation ($p < 0.001$) were found between toluene, m-xylene, ethylbenzene, o-xylene, p-xylene and total BTEX exposure levels in the breathing zone (Table 3): in fact, emissions and combustion products from gasoline vehicles include a large variety of chemicals and additives, *e.g.* toluene; when those compounds are emitted from the same source, they have a nearly constant emission ratio (Barbieri et al., 2008).

3.3. Biomarkers of exposure

3.3.1. Benzene

The median level of post-shift *t,t*-MA in traffic policemen was 13-fold higher than that of office policemen ($p < 0.001$) ($p = p =$ (Table 4). The concentrations of pre- and post-shift *t,t*-MA were significantly different in office policemen ($p = 0.03$). The concentration of post-shift S-PMA in traffic policemen was not shown to be significantly different from that of the office policemen. No significant associations were found between personal benzene, toluene, ethylbenzene, m-xylene, o-xylene, p-xylene or total BTEX exposure and S-PMA or *t,t*-MA levels (Table 3).

3.3.2. 1,3-Butadiene

The median level of pre-shift MHBMA in traffic policemen was 6.5-fold higher than that of office policemen ($p < 0.001$). The concentration of post-shift MHBMA in traffic policemen was not shown to be significantly different from that of the office policemen (Table 4). The median level of pre-shift DHBMA in traffic policemen was 2.7-fold higher than that of office policemen ($p < 0.001$) and 2.8-fold higher at post-shift ($p < 0.01$) compared between traffic and office policemen. A strong significant correlation between personal toluene, ethylbenzene, m-xylene, o-xylene, p-xylene or total BTEX exposure and urinary post-shift DHBMA (Table 3). The association between personal total BTEX exposure and post-shift DHBMA may indicate the same source of emissions, *i.e.* automobile exhaust. Significant difference was found between sub-groups of non-smokers, light to moderate smokers and heavy smokers only at pre-shift *t,t*-MA levels ($p = 0.01$) (Kruskal-Wallis test). Median levels of urinary post-shift MHBMA in the urine of heavy smokers was 5.6-fold higher than in non-smokers ($p = 0.025$) (Mann-Whitney test). Urinary S-PMA and DHBMA values did not discriminate exposure resulting from smoking habits and no significant difference was found between smokers and non-smokers (data not shown). Nevertheless, significant difference was found between sub-groups of non-smokers, light to moderate smokers and heavy smokers at pre-shift and post-shift MHBMA results ($p = 0.007$ and $p = 0.003$) (Kruskal-Wallis test). Median levels of urinary post-shift MHBMA in the urine of light to moderate smokers and heavy smokers were, respectively, 4.9-fold and 4.1-fold higher than in non-smokers.

3.4. Multiple regression analysis

Based on multiple regression analyses, smoking consumption was not found to have a significant influence on DHBMA which was strongly related to logBTEX ($\beta = 0.535$; $p <$

277 0.001), showing that there is a difference in sensitivity between the two urinary metabolites
278 of BD exposure measured in this study. Nevertheless, age was negatively related to
279 logDHBMA ($\beta = -0.286$; $p = 0.045$).

280

4. DISCUSSION

Environmental exposure to VOCs among workers has been already described elsewhere (Barbieri et al., 2008; Pilidis et al., 2008; Arayasiri et al., 2010; Manini et al., 2010), whereas it has not been reported yet for Beirut traffic policemen. For these reasons, we have undertaken an extensive investigation of the occupational exposure to VOCs, notably, benzene and BD, which are carcinogenic substances in urban air from incomplete combustion of fossil fuels, through personal air monitoring, as well as through the use of biomarkers of exposure. In a previous studies conducted in Beirut, it has been reported that the spatial distribution of VOCs emissions are mostly over Beirut and its suburbs, where there are dense populations and heavy traffic (Waked et al., 2012). While ambient air monitoring of VOCs-compounds has not been reported at roadsides in Beirut area, individual exposure levels to BTEX compounds in traffic policemen were significantly higher than in office policemen. These levels were far below the Occupational Safety and Health Administration (OSHA) exposure limits, 8-h time weight average of 3.2 mg m^{-3} for benzene (OSHA, 1999), of 375 mg m^{-3} for toluene (OSHA, 1999), of 435 mg m^{-3} for ethylbenzene (OSHA, 1999) and of 435 mg m^{-3} for xylene (OSHA, 2008). Moreover, our results have shown that each exposure to one vehicle increases the level of individual exposure to benzene, o-xylene and to p-xylene due to the fact that xylene is primarily released from motor vehicle exhaust where it is used as a solvent (Kandyala et al., 2010). Evidence presented herein extended these observations by showing that these personal exposure levels indicated higher emissions related to vehicles in Beirut area that are highly traffic congested and to the fact that in Lebanon personal vehicles are the prevailing mode of transport.

In addition, individual exposure to benzene (mean = $48.8 \text{ } \mu\text{g m}^{-3}$) in the traffic policemen in this study was higher than that measured in traffic policemen in Bologna/Italy (mean = 17.3

306 $\mu\text{g m}^{-3}$) (Barbieri et al., 2008), in Ioannina/Greece (mean = $30 \mu\text{g m}^{-3}$) (Pilidis et al., 2008), in
307 Prague/Czech Republic (mean concentration in February = 6.99 and in May = $4.53 \mu\text{g m}^{-3}$)
308 (Rossnerova et al., 2009) and in Bangkok/Thailand (mean= $38.24 \mu\text{g m}^{-3}$) (Arayasiri et al.,
309 2010). The same for other BTEX compounds, individual exposure to toluene, ethylbenzene,
310 m-xylene and o-xylene in the traffic policemen in this study were strongly higher than that
311 measured in traffic policemen in Prague/Czech Republic (mean concentration in May =
312 13.99, 3.25, 10.95 and $3.75 \mu\text{g m}^{-3}$, respectively) (Rossnerova et al., 2009). Difference in
313 many of environment factors, such as traffic characteristics, quality of fuel, meteorological
314 conditions, building characteristics of the area, and difference in physical activity in the
315 workplace may contribute to the difference in the levels of individual VOCs exposure.
316 Although excellent spearman correlations ($p < 0.001$) were found between toluene,
317 ethylbenzene, m-xylene, o-xylene, p-xylene and total BTEX exposure levels, we did not find
318 significant correlation between benzene and none of BTEX compounds. Even if these
319 compounds were emitted from the same source (Barbieri et al., 2008), we suggest that lack of
320 any such correlation may be due to the fact that some individual values reported for benzene
321 exposure were below the LOD ($< 0.3 \mu\text{g m}^{-3}$).

322 In our study, *t,t*-MA concentration was found to be a better indicator of the levels of
323 benzene exposure than S-PMA. Urinary post-shift *t,t*-MA levels were significantly higher in
324 traffic policemen compared with office policemen (Table 4). In a previous studies with Italian
325 policemen who had levels of individual benzene exposure of $17.3 \mu\text{g m}^{-3}$ (Barbieri et al.,
326 2008), *i.e.* lower than in this study, post-shift *t,t*-MA levels were comparable ($44 \mu\text{g m}^{-3}$ for
327 non-smokers and $120 \mu\text{g m}^{-3}$ for smokers) (Manini et al., 2010), while the post-shift S-PMA
328 levels were much lower in both studies (approximately 7-fold). However, the levels of S-
329 PMA and *t,t*-MA were not significantly different between pre- and post-shift samples,
330 excepted for *t,t*-MA results within office policemen group ($p = 0.03$). A close examination at

individual results reveals a decrease of urinary S-PMA by the end of the work shift. A previous study, conducted by Qu and colleagues (2000), estimated the urinary half life of elimination ($t_{1/2}$) of S-PMA to be 12.8 h. Thus, higher levels in pre-shift S-PMA could be the result of cumulative exposure to benzene during the previous day. Moreover, ingestion of unknown amounts of dietary sorbic acid and glutathione-S-transferase (*GSTM1*, *GSTT1* and *GSTA1*) polymorphism may influence variability in metabolites levels (Manini et al., 2010). Moreover, we denote some urinary S-PMA and *t,t*-MA values below the LOQ ($< 1 \mu\text{g L}^{-1}$ and $< 10 \mu\text{g L}^{-1}$, respectively). It seems that biotransformation of benzene to its metabolites is reduced by co-exposure to other chemicals and additives emitted from gasoline vehicles (Barbieri et al., 2008) such as toluene (a competitive inhibitor of benzene for CYP metabolism) which would affect the pharmacokinetics and metabolism of benzene, including conjugation with glutathione and subsequent mercapturic acid excretion (Johnson et al., 2007). Accordingly to previous studies, no significant associations were found between personal benzene or personal BTEX compounds exposure and S-PMA or *t,t*-MA levels (Table 3) (Barbieri et al., 2008; Arayasiri et al., 2010). The lack of any such correlation may be due to the confounding effect of smoking on metabolite excretion (Carrieri et al., 2010; 2012) or to the fact that some values reported for benzene exposure, urinary PMA and *t,t*-MA were below the LOQ. Moreover, urinary *t,t*-MA level may be affected by the diet of the subjects examined (Carrieri et al., 2010; Weisel, 2010). For BD exposure in ambient air, a limited number of biomonitoring studies have been conducted. Pre- and post-shift DHBMA levels in traffic policemen were significantly higher than in office policemen (Table 4). We found that urinary pre-shift MHBMA and DHBMA concentrations in traffic policemen were significantly higher than office policemen, which indicates that elimination of those metabolites from previous BD exposure is not yet completed. This suggests that the average urinary $t_{1/2}$ of MHBMA and DHBMA is somewhat protracted compared to the $t_{1/2}$ of other

mercapturic acids, such as the benzene metabolite S-PMA and *t,t*-MA, which have an average $t_{1/2}$ of 12.8 h and 13.7 h, respectively (Qu et al., 2000; van Sittert et al., 2000; Albertini et al., 2003). A close examination at individual results reveals a decrease of urinary MHBMA level by the end of the work shift. In view of the complexity of the metabolic pathways involved in the biotransformation of butadiene, we expected an effect of the genetic polymorphisms in xenobiotic metabolizing enzymes which affects the excretion of urinary metabolites. We also suggest that co-exposure to aromatic hydrocarbons (*e.g.* toluene) may cause a decrease in epoxide metabolites levels, namely formation via cytochrome P-450 2E1, and a decreased excretion urinary metabolites of BD (Johnson et al., 2007; Vacek et al., 2010; Fustinoni et al., 2012).

Alternatively, DHBMA concentrations were found to be a better indicator of the levels of BD exposure in urban air than MHBMA, as reported previously (Sapkota et al., 2006). Pre- and post-shift DHMBA levels could significantly distinguish between office and traffic policemen and showed a better correlation with personal total BTEX exposure.

Multiple regression analysis applied to evaluate the contribution of each predictor to the variability of urinary biomarkers of benzene and BD. Traffic and office policemen stated to have smoked during working. Although levels of BD in the breathing zone of study subjects have not been reported, it seems that smoking habits influence the total intake of BD (WHO, 2000). Unexpectedly, there was not a marked effect of smoking habits on DHBMA. Multiple regression analysis showed that the levels of individual BTEX exposure significantly contributed to increase urinary DHBMA ($\beta=0.535$, $p < 0.001$). Accordingly, DHBMA values did not discriminate exposure resulting from smoking habits and no significant difference was found between smokers and non-smokers.

In conclusion, these results indicated that traffic policemen, who are exposed to benzene and BD at the roadside in central Beirut, are potentially at a higher risk for development of

381 diseases such as cancer than office policemen. Further studies need to focus on the lifestyle
382 and genetic factors that may affect the background levels of S-PMA, *t,t*-MA, MHBMA and
383 DHBMA. Given chaotic traffic conditions, quality of fuel, high rate of passenger cars
384 ownership, meteorological conditions, building characteristics of the area, and high pollution
385 rates observed in Beirut, increased urinary exposure biomarker would result in a higher risk
386 in the exposed subjects. Accordingly, public authorities should particularly set policies aimed
387 to reduce traffic-related air pollution.

388

389 Funding

390 This research was funded by Agence Universitaire de la Francophonie (PCSI;
391 reference: 6317PS011)

392

393 Acknowledgments

394 The authors express their sincere thanks to the Maj. Gen. Ashraf Rifi, General
395 Director of the Lebanese Internal Security Forces for his encouragement, and to the Major
396 Elie Kallas and the Major Hanna Laham for providing facilities during the study. The authors
397 declare that there is no conflict of interest.

398

399 **References**

- 400 Abbas, I., Garçon, G., Saint-Georges, F., Andre, V., Gosset, P., Billet, S., Goff, J.L., Verdin,
401 A., Mulliez, P., Sichel, F., Shirali, P., 2013. Polycyclic aromatic hydrocarbons within
402 airborne particulate matter (PM_{2.5}) produced DNA bulky stable adducts in a human lung
403 cell coculture model. *J Appl Toxicol.* 33:109-19.
- 404 Albertini, R.J., Sram, R.J., Vacek, P.M., Lynch, J., Wright, M., Nicklas, J.A., Boogaard, P.J.,
405 Henderson, R.F., Swenberg, J.A., Tate, A.D., Ward, J.B., Jr., 2001. Biomarkers for
406 assessing occupational exposures to 1,3-butadiene. *Chem Biol Interact.* 135:429–453.
- 407 Arayasiri, M., Mahidol, C., Navasumrit, P., Autrup, H., Ruchirawat, M., 2010. Biomonitoring
408 of benzene and 1,3-butadiene exposure and early biological effects in traffic policemen.
409 *Sci Total Environ.* 408:4855-62.
- 410 Avogbe, P., Ayi-Fanou, L., Cachon, B., Chabi, N., Delbende, A., Dewaele, D., Aissi, F.,
411 Cazier, F., and Sanni A., 2011. Hematological changes among Beninese motor-bike taxi
412 drivers exposed to benzene by urban air pollution. *Afr. J. Environ. Sc. Technol.* 5: 464-
413 472
- 414 Barbieri, A., Violante, F.S., Sabatini, L., Graziosi, F., Mattioli, S., 2008. Urinary biomarkers
415 and low-level environmental benzene concentration: assessing occupational and general
416 exposure. *Chemosphere.* 74:64-9.
- 417 Billet, S., Paget, V., Garçon, G., Heutte, N., André, V., Shirali, P., Sichel, F., 2010. Benzene-
418 induced mutational pattern in the tumour suppressor gene TP53 analysed by use of a
419 functional assay, the functional analysis of separated alleles in yeast, in human lung
420 cells. *Arch Toxicol.* 84:99-107.
- 421 Carugno, M., Pesatori, A.C., Dioni, L., Hoxha, M., Bollati, V., Albetti, B., Byun, H.M.,
422 Bonzini, M., Fustinoni, S., Cocco, P., Satta, G., Zucca, M., Merlo, D.F., Cipolla, M.,
423 Bertazzi, P.A., Baccarelli, A., 2012. Increased mitochondrial DNA copy number in

424 occupations associated with low-dose benzene exposure. *Environ Health Perspect.*
 425 120:210-5.

426 Cheng, X., Zhang, T., Zhao, J., Zhou, J., Shao, H., Zhou, Z., Kong, F., Feng, N., Sun, Y.,
 427 Shan, B., Xia, Z., 2012. The association between genetic damage in peripheral blood
 428 lymphocytes and polymorphisms of three glutathione S-transferases in Chinese workers
 429 exposed to 1,3-butadiene. *Mutat Res.* 750:139-46.

430 Clifford P. Weisel, 2010. Benzene exposure: An overview of monitoring methods and their
 431 findings. *Chem-Biol Interact.* 184:58–66

432 Dagher, Z., Garçon, G., Billet, S., Gosset, P., Ledoux, F., Courcot, D., Aboukais, A., Shirali,
 433 P., 2006. Activation of different pathways of apoptosis by air pollution particulate matter
 434 (PM_{2.5}) in human epithelial lung cells (L132) in culture. *Toxicol.* 225:12-24.

435 EU-RAR, 2002. European Union Risk Assessment Report 1,3-Butadiene.

436 Fustinoni, S., Campo, L., Satta, G., Campagna, M., Ibba, A., Tocco, M., Atzeri, S., Avataneo,
 437 G., Flore, C., Meloni, M., Bertazzi, P., Cocco, P., 2012. Environmental and lifestyle
 438 factors affect benzene uptake biomonitoring of residents near a petrochemical plant.
 439 *Environ Int.* 39: 2-7.

440 Fustinoni, S., Consonni, D., Campo, L., Buratti, M., Colombi, A., Pesatori, A.C., Bonzini,
 441 M., Bertazzi, P.A., Foà, V., Garte, S., Farmer, P.B., Levy, L.S., Pala, M., Valerio, F.,
 442 Fontana, V., Desideri, A., Merlo, D., 2005. Monitoring low benzene exposure:
 443 comparative evaluation of urinary biomarkers, influence of cigarette smoking, and genetic
 444 polymorphisms. *Cancer Epidemiol. Biomarkers Prev.* 14, 2237–2244.

445 Han, X., Naeher, L., 2006. A review of traffic-related air pollution exposure assessment
 446 studies in the developing world. *Environ Int.* 32:106-20

447 IARC, 2008. Monographs On The Evaluation Of The Carcinogenic Risk Of Chemicals To
 448 Humans. Volume 97. 1,3-Butadiene, Ethylene Oxide and Vinyl Halides
 449 (<http://monographs.iarc.fr>).
 450 IARC, 2009. A Review Of Human Carcinogens, Part F: Chemical Agents And Related
 451 Occupations. *Lancet Oncol* 10:1143-1144.
 452 Ibrahim, K.S., Amer, N.M., El-Dossuky, E.A., Emara, A.M., El-Fattah, A.E., Shahy, E.M.,
 453 2012. Hematological effects of benzene exposure with emphasis on muconic acid as
 454 biomarker in exposed workers. *Toxicol Ind Health*. 0748233712458141.
 455 Johnson, E.S., Langård, S., Lin, Y.S., 2007. A critique of benzene exposure in the general
 456 population. *Sci Total Environ*. 374:183-98.
 457 Kamal, A., Malik, R.N., Fatima, N., Rashid, A., 2012. Chemical exposure in occupational
 458 settings and related health risks: a neglected area of research in Pakistan. *Environ*
 459 *Toxicol Pharmacol*. 34:46-58.
 460 Kandyala, R., Raghavendra, S.P., Rajasekharan, S.T., 2010. Xylene: An overview of its
 461 health hazards and preventive measures. *J Oral Maxillofac Pathol*. 14:1–5.
 462 Knibbs, L.D., Morawska, L., 2012. Traffic-related fine and ultrafine particle exposures of
 463 professional drivers and illness: An opportunity to better link exposure science and
 464 epidemiology to address an occupational hazard? *Environ Int*. 49:110-4.
 465 Koturbash, I., Scherhag, A., Sorrentino, J., Sexton, K., Bodnar, W., Tryndyak, V.,
 466 Latendresse, J.R., Swenberg, J.A., Beland, F.A., Pogribny, I.P., Rusyn, I., 2011.
 467 Epigenetic alterations in liver of C57BL/6J mice after short-term inhalational exposure
 468 to 1,3-butadiene. *Environ Health Perspect*. 119:635-40.
 469 Maffei, F., Hrelia, P., Angelini, S., Carbone, F., Forti, G. V., Barbieri, A., 2005. Effects of
 470 environmental benzene: Micronucleus frequencies and haematological values in traffic
 471 police working in an urban area. *Mutat. Res*. 583 :1–11.

472 Manini, P., De Palma, G., Andreoli, R., Mozzoni, P., Poli, D., Goldoni, M., Petyx, M.,
 473 Apostoli, P., Mutti, A., 2010. Occupational exposure to low levels of benzene:
 474 Biomarkers of exposure and nucleic acid oxidation and their modulation by polymorphic
 475 xenobiotic metabolizing enzymes. *Toxicol Lett.* 193:229-35.

476 Occupational Safety and Health Administration (OSHA), US. Department of Labor. OSHA
 477 standard -29 CFR 1910.1028 (7/1/1999), from National Library of Medicine, HSDB:
 478 Benzene, CASRN: 71-43-2; 1999 ([http://toxnet.nlm.nih.gov/cgi-](http://toxnet.nlm.nih.gov/cgi-bin/sis/search/f?./temp/~F3eDcl:1)
 479 [bin/sis/search/f?./temp/~F3eDcl:1](http://toxnet.nlm.nih.gov/cgi-bin/sis/search/f?./temp/~F3eDcl:1) Last accessed 07/12/2012).

480 OSHA, OSHA standard -29 CFR 1910.1000 (2/10/2008), from U.S. National Archives and
 481 Records Administration's Electronic Code of Federal Regulations, HSDB: Xylene,
 482 CASRN: 1330-20-7; 2008 (Last accessed 07/12/2012).

483 OSHA, OSHA standard -29 CFR 1910.1000 (7/1/1999), from National Library of Medicine,
 484 HSDB: Toluene, CASRN: 108-88-3; 1999 (Last accessed 07/12/2012).

485 OSHA, OSHA standard -29 CFR 1910.1000 (7/1/1999), from National Library of Medicine,
 486 HSDB: Ethylbenzene, CASRN: 100-41-4; 1999 (Last accessed 07/12/2012).

487 Pamela M. Vacek, Richard J. Albertini, Radim J. Sram, Patricia Upton, James A. Swenberg,
 488 Hemoglobin adducts in 1,3-butadiene exposed Czech workers: Female–male
 489 comparisons, *Chemico-Biological Interactions*, Volume 188, Issue 3, 5 December 2010,
 490 Pages 668-676.

491 Peluso, M., Bollati, V., Munnia, A., Srivatanakul, P., Jedpiyawongse, A., Sangrajang, S.,
 492 Piro, S., Ceppi, M., Bertazzi, P.A., Boffetta, P., Baccarelli, A.A., 2012. DNA
 493 methylation differences in exposed workers and nearby residents of the Ma Ta Phut
 494 industrial estate, Rayong, Thailand. *Int J Epidemiol.* 41:1753-60.

495 Pilidis, G.A., Karakitsios, S.P., Kassomenos, P.A., Kazos, E.A., Stalikas, C.D., 2009.
 496 Measurements of benzene and formaldehyde in a medium sized urban environment.

497 Indoor/outdoor health risk implications on special population groups. *Environ Monit*
 498 *Assess.* 150:285-94.

499 Qu, Q., Melikian, A.A., Li, G., Shore, R., Chen, L., Cohen, B., Yin, S., Kagan, M.R., Li, H.,
 500 Meng, M., Jin, X., Winnik, W., Li, Y., Mu, R., Li, K., 2000. Validation of biomarkers in
 501 humans exposed to benzene: urine metabolites. *Am J Ind Med.* 37:522-31.

502 Rekhadevi, P.V., Rahman, M.F., Mahboob, M., Grover, P., 2010. Genotoxicity in filling
 503 station attendants exposed to petroleum hydrocarbons. *Ann Occup Hyg.* 54:944-54.

504 Rossnerova, A., Spatova, M., Rossner, P., Solansky, I., Sram, R.J., 2009. The impact of air
 505 pollution on the levels of micronuclei measured by automated image analysis. *Mutat*
 506 *Res.* 669:42-7.

507 Ruchirawat, M., Navasumrit, P., Settachan, D., 2010. Exposure to benzene in various
 508 susceptible populations: co-exposures to 1,3-butadiene and PAHs and implications for
 509 carcinogenic risk. *Chem Biol Interact.* 184:67-76.

510 Sangaraju, D., Goggin, M., Walker, V., Swenberg, J., Tretyakova, N., 2012. NanoHPLC-
 511 nanoESI(+)-MS/MS quantitation of bis-N7-guanine DNA-DNA cross-links in tissues of
 512 B6C3F1 mice exposed to subppm levels of 1,3-butadiene. *Anal Chem.* 84:1732-9.

513 Sapkota, A., Halden, R.U., Dominici, F., Groopman, J.D., Buckley, T.J., 2006. Urinary
 514 biomarkers of 1,3-butadiene in environmental settings using liquid chromatography
 515 isotope dilution tandem mass spectrometry. *Chem Biol Interact.* 160:70-9.

516 Seow, W.J., Pesatori, A.C., Dimont, E., Farmer, P.B., Albeti, B., 2012. Urinary Benzene
 517 Biomarkers and DNA Methylation in Bulgarian Petrochemical Workers: Study Findings
 518 and Comparison of Linear and Beta Regression Models. *PLoS One* 7:e50471.

519

520 van Sittert, N., Megens, H., Watson, W., Boogaard, P., 2000. Biomarkers of exposure to 1,3-
 521 butadiene as a basis for cancer risk assessment. *Toxicol Sci.* 56:189-202. Tabish, A.M.,

522 Poels, K., Hoet, P., Godderis, L., 2012. Epigenetic factors in cancer risk: effect of
 523 chemical carcinogens on global DNA methylation pattern in human TK6 cells. PLoS
 524 One. 7:e34674.

525 Waked, A., Afif, C., Seigneur, C., 2012. An atmospheric emission inventory of
 526 anthropogenic and biogenic sources for Lebanon. Atmos. Environ. 50,88-96

527 WHO, 1996. Biological Monitoring Of Chemical Exposure In The Workplace, vol. 1,
 528 Geneva: World Health Organization.

529 WHO, 2000. Butadiene Air Quality Guidelines. Chapter 5.3 - Second Edition. Regional
 530 Office for Europe, Copenhagen, Denmark.

531 Xiang, M., Ao, L., Yang, H., Liu, W., Sun, L., Han, X., Li, D., Cui, Z., Zhou, N., Liu, J., Cao,
 532 J., 2012. Chromosomal damage and polymorphisms of metabolic genes among 1, 3-
 533 butadiene-exposed workers in a matched study in China. Mutagenesis. 27:415-21.

534

535 TABLES

536

537 **Table 1.** Summary of information collected by questionnaire.

	Traffic policemen (n=24)	Office policemen (n=23)	<i>p</i> value
Age (years, mean $\pm$ SD)	27.4 $\pm$ 4.6	29.1 $\pm$ 3.6	0.06
Percentage of never/current-smokers	37	56	0.19
Cigarettes/d (mean $\pm$ SD)	21 $\pm$ 18	26.2 $\pm$ 23	0.633
Sampling period (h, mean $\pm$ SD)	33.25 $\pm$ 3.1	45.2 $\pm$ 3.4	<0.001
BMI (kg m ⁻² , mean $\pm$ SD)	25.6 $\pm$ 4.6	27.9 $\pm$ 4.3	0.125

538 Significant at $p < 0.05$. Student's *t* test.

539 Table 2. Distribution of individual VOCs exposure in subgroups of policemen. Values are expressed as geometric medians, means and
540 geometric standard deviations. Concentrations are expressed as $\mu\text{g}/\text{m}^3$.

Parameter	Traffic policemen (n=24)			Parameter	Office policemen (n=23)			<i>p value</i>
	Median	Min-Max	Mean $\pm$ SD		Median	Min-Max	Mean $\pm$ SD	
Benzene (n=21)	0.3	0.3-867.8	48.8 $\pm$ 189.3	Benzene (n=22)	0.3	0.3-10.8	1.2 $\pm$ 2.3	n.s.
Toluene (n=20)	101.8	29.8-9788.4	11520. $\pm$ 2393.0	Toluene (n=22)	9.3	0.44-46.3	12.1 $\pm$ 11.3	<0.001
Ethylbenzene (n=21)	58.1	6.7-5845.2	662.3 $\pm$ 1420.3	Ethylbenzene (n=22)	4.5	0.88-19.1	5.7 $\pm$ 4.4	<0.001
mXylene (n=21)	102.8	29.1-1209.6	196.7 $\pm$ 287.6	mXylene (n=22)	2.0	0.65-10.6	2.5 $\pm$ 2.2	<0.001
oXylene (n=21)	21.3	1.0-1839.1	177.7 $\pm$ 432.3	oXylene (n=22)	1.7	0.58-5.64	2.0 $\pm$ 1.3	<0.001
pXylene (n=15)	3.9	1.2-30.1	9.6 $\pm$ 10.0	pXylene (n=20)	0.4	0.18-2.1	0.6 $\pm$ 0.4	<0.001
BTEX (n=21)	295.1	85.8-18709.1	2189.0 $\pm$ 4450.8	BTEX (n=22)	21.4	3.6-84.1	24.3 $\pm$ 19.6	<0.001

541 Significant at $p < 0.05$. Mann-Whitney test, n.s. = not significant.

542

543 Table 3. Spearman correlation coefficients between different variables.

	Tol	Ethylbz	mXyl	oXyl	pXyl	BTEX	S-PMA ¹	S-PMA ²	<i>t,t</i> -MA ¹	<i>t,t</i> -MA ²	MHBMA ¹	MHBMA ²	DHBMA ¹	DHBMA ²
Bz	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Tol		0.962 ^{**}	0.982 ^{**}	0.939 ^{**}	0.933 ^{**}	0.989 ^{**}	0.358 [*]	-	-	-	-	-	0.505 ^{**}	0.610 ^{**}
Ethylbz			0.965 ^{**}	0.931 ^{**}	0.950 ^{**}	0.975 ^{**}	0.354 [*]	-	-	-	0.317 [*]	-	0.496 ^{**}	0.639 ^{**}
mXyl				0.951 ^{**}	0.944 ^{**}	0.988 ^{**}	0.381 [*]	-	-	-	0.357 [*]	-	0.490 ^{**}	0.607 ^{**}
oXyl					0.946 ^{**}	0.944 ^{**}	0.394 ^{**}	-	-	-	-	-	0.411 ^{**}	0.565 ^{**}
pXyl						0.943 ^{**}	0.398 [*]	-	-	-	-	-	0.370 [*]	0.597 ^{**}
BTEX							0.401 ^{**}	-	-	-	0.345 [*]	-	0.495 ^{**}	0.616 ^{**}
S-PMA ¹								0.467 ^{**}	0.464 ^{**}	-	0.389 ^{**}	-	0.384 ^{**}	0.356 [*]
S-PMA ²									-	-	-	0.374 [*]	-	0.320 [*]
<i>t,t</i> -MA ¹										-	0.385 ^{**}	-	-	-
<i>t,t</i> -MA ²											-	-	-	-
MHBMA ¹												0.352 [*]	0.538 ^{**}	0.540 ^{**}
MHBMA ²													-	-
DHBMA ¹														0.716 ^{**}

544 Missing values correspond to lack of correlation between parameters. Only significant data are represented. ¹ Pre-shift; ² Post-shift. ** $p < 0.001$;
545 * $p < 0.05$

546 Table 4. Biomarkers of exposure in traffic and office policemen.

547

Parameter	Study groups	
	Traffic policemen	Office policemen
Urinary <i>t,t</i> -MA (µg/g creatinine)		
Pre-shift	64.7±141.5	14±13.3
	8.2 (3-586.7)	11.2 (2.5-56.5)
	(n=24)	(n=23)
Post-shift	84.4±100.6 ^a	55.5±87 [*]
	61 (2.4-444.8)	18.3 (2.5-343.4)
	(n=24)	(n=22)
Urinary S-PMA (µg/g creatinine)		
Pre-shift	6.2±6 ^a	2.6±2
	4 (1.1-25)	1.7 (0.9-8.1)
	(n=24)	(n=23)
Post-shift	5.3±9	2.7±2.4
	2.6 (0.5-45.5)	1.9 (0.4-10.2)
	(n=24)	(n=22)
Urinary MHBMA (µg/g creatinine)		
Pre-shift	24.7±23.3 ^a	17.7±38.7
	20.1 (2.4-108.3)	3.1 (1.3-147.6)
	(n=24)	(n=23)
Post-shift	18.7±20.1	18.8±31
	10.2 (0.2-88.6)	7.5 (1.2-120)
	(n=24)	(n=22)

Urinary DHBMA (µg/g creatinine)

Pre-shift	180.4±73.8 ^a	69.2±43.6
	182.9 (47.6-333.3)	66.5 (13.2-239.3)
	(n=24)	(n=23)
Post-shift	207.5±112.2 ^b	73.3±45.3
	188.6 (80-588.6)	67.4 (23.9-186.7)
	(n=23)	(n=22)

Urinary Cotinine (µg/g creatinine)

Pre-shift	767.8±1138.5 ^a	188.6±339.2
	486.9 (16.1-5240)	24.6 (8.5-1200.8)
	(n=24)	(n=23)
Post-shift	601.0±780.6	205.5±393.2
	459.3 (12.1-3583.9)	19.9 (7.9-1362.4)
	(n=24)	(n=22)

548 Values are expressed as mean±SD on the first line and median (min-max) on the second line

549 of each parameter.

550 Statistically significant difference from office policemen at ^a $p < 0.001$ or ^b $p < 0.01$.

551 * Statistically significant difference from the corresponding pre-shift at $p = 0.03$.